



Content

Acknowledgements	I
List of figures	VII
List of tables	XIII
Abbreviations	XV
1 Introduction and Intention	1
2 General	3
2.1 Polyphenols.....	3
2.1.1 Classification of Polyphenols	3
2.1.1.1 Non-flavonoids	4
2.1.1.1.1 Hydroxybenzoic acids.....	4
2.1.1.1.2 Hydroxycinnamic acids.....	4
2.1.1.2 Flavonoids.....	5
2.1.1.2.1 Flavanones	6
2.1.1.2.2 Flavones	6
2.1.1.2.3 Flavonols.....	7
2.1.1.2.4 Flavan-3-ols: Catechins and Proanthocyanidins	8
2.1.1.2.5 Anthocyanidins: Anthocyanins	10
2.1.2 Biosynthesis of phenolics, polyphenols and anthocyanins	18
2.1.3 Absorption, Bioactivity and Metabolism of anthocyanins	27
2.1.3.1 Anthocyanin absorption and metabolism.....	27
2.1.3.2 Bioactivity of Anthocyanins	33
2.1.3.2.1 Antioxidant activity	33
2.1.3.2.2 Anti-carcinogenic effects	35
2.1.3.2.3 Anti-inflammatory effects.....	37
2.1.3.2.4 Anti-cardiovascular effects	38
2.1.3.2.5 Antidiabetic effects	38
2.1.3.2.6 Neuroprotective effects	39
2.1.3.2.7 Antibacterial, antimicrobial and antiviral activity	40
2.1.3.2.8 Gastro protective effects	40



Content

2.1.3.2.9	Skin protective effect	41
2.1.3.2.10	Effect on vision	41
2.2	Raw materials	42
2.2.1	Blackberry (<i>Rubus fruticosus</i>)	42
2.2.2	Black chokeberry (<i>Aronia melanocarpa</i>)	43
2.2.3	Sour cherries (<i>Prunus cerasus</i> L.)	44
2.2.4	Black carrots (<i>Daucus carota</i> ssp. <i>sativus</i> var. <i>atrorubens</i> Alef.)	45
2.2.5	Purple sweet potato (<i>Ipomoea batatas</i> L.)	46
2.2.6	Roselle (<i>Hibiscus sabdariffa</i> L.)	47
2.2.7	Butterfly pea (<i>Clitoria ternatea</i> L.)	48
2.3	Preparative techniques for separation	49
2.3.1	Countercurrent Chromatography	49
2.3.1.1	High-Speed Countercurrent Chromatography (HSCCC)	52
2.3.1.2	High Performance Countercurrent Chromatography (HPCCC)	53
2.3.1.3	Low-Speed Rotary Countercurrent Chromatography (LSRCCC)	54
2.3.2	Membrane chromatography	55
3	Results and Discussion	57
3.1	Working outline	57
3.2	Blackberry	58
3.2.1	Chemical composition	58
3.2.2	Preparative separations	63
3.2.2.1	Membrane chromatography	63
3.2.2.2	Countercurrent Chromatography	66
3.3	Black chokeberry	71
3.3.1	Chemical composition	71
3.3.2	Preparative separations	77
3.3.2.1	Membrane chromatography	77
3.3.2.2	Countercurrent Chromatography	80
3.4	Sour cherries	85
3.4.1	Chemical composition	85



Content

3.4.2	Preparative separations	90
3.4.2.1	Membrane chromatography	90
3.5	Black carrots	94
3.5.1	Chemical composition	94
3.5.2	Preparative separations	98
3.5.2.1	Membrane chromatography	98
3.5.2.2	Countercurrent Chromatography	101
3.6	Purple sweet potato	105
3.6.1	Chemical composition	105
3.6.2	Preparative separations	108
3.6.2.1	Membrane chromatography	108
3.6.2.2	Countercurrent Chromatography	111
3.7	Roselle	114
3.7.1	Chemical composition	114
3.7.2	Preparative separations	117
3.7.2.1	Membrane chromatography	117
3.7.2.2	Countercurrent Chromatography	122
3.8	Butterfly pea	124
3.8.1	Chemical composition	124
3.8.2	Preparative separations	129
3.8.2.1	Membrane chromatography	129
3.8.2.2	Countercurrent Chromatography	131
3.9	Authenticity and Food Adulterations	133
4	Material and Methods	135
4.1	Plant materials	135
4.2	Chemicals	137
4.3	Extraction of polyphenols	138
4.4	Adsorption chromatography on Amberlite XAD-7	138
4.5	Solvent precipitation	138
4.6	Device parameters	139



Content

4.6.1	HPLC-DAD	139
4.6.1.1	Analytical HPLC-DAD.....	139
4.6.1.2	Preparative HPLC-UV/VIS	141
4.6.2	HPLC-DAD-ESI-MS ⁿ	142
4.6.3	TLC.....	143
4.6.4	UV Spectrophotometer	143
4.6.4.1	Total phenolic content by Folin Ciocalteu method.....	144
4.6.5	Membrane chromatography.....	144
4.6.6	Countercurrent Chromatography.....	145
4.6.6.1	High-Speed Countercurrent Chromatography	145
4.6.6.2	High Performance Countercurrent Chromatography.....	145
4.6.6.3	Low-Speed Rotary Countercurrent Chromatography	146
4.6.7	Nuclear Magnetic Resonance Spectroscopy.....	148
5	Summary and Conclusion	155
6	Zusammenfassung.....	157
7	References	159



List of figures

Fig. 2.1. Classes of polyphenols (Iriti and Faoro, 2009).....	3
Fig. 2.2. Chemical structure of hydroxybenzoic acids.	4
Fig. 2.3. Chemical structure of hydroxycinnamic acids.	4
Fig. 2.4. Structures of principal flavonoids subgroups (Crozier et al., 2009; Iriti and Faoro, 2009).	5
Fig. 2.5. Chemical structure of flavanones and some examples.	6
Fig. 2.6. Structure of flavones.	7
Fig. 2.7. Chemical structures of flavonol aglycones.	7
Fig. 2.8. Chemical structure of flavan-3-ols.	8
Fig. 2.9. Some chemical structures of A-type and B-type proanthocyanidins.....	9
Fig. 2.10. Chemical structure of sugars, which usually occur in anthocyanins.	12
Fig. 2.11. Structure of aliphatic and aromatic acids that can exist in anthocyanins.	13
Fig. 2.12. Equilibrium forms of anthocyanidins. R = glycoside; R ₁ and R ₂ can be OH, CH ₃ or OCH ₃ groups, according to the type of aglycone.....	14
Fig. 2.13. Specific positions of anthocyanidins to react (Mateus and Freitas, 2009; Andersen and Jordheim, 2013).	15
Fig. 2.14. Reaction on the C-Ring, generating hemiketals and sulfites. R = glycoside; R ₁ and R ₂ can be OH, CH ₃ or OCH ₃ groups, according to the type of aglycone.....	16
Fig. 2.15. The formation of pyranoanthocyanins: reaction between Mv-3-glc and carbonyl compounds.	17
Fig. 2.16. Metalloanthocyanin: Representation of interaction between cation and anthocyanin in commelinin (left), self-assembly of metalloanthocyanins (middle) and protocyanidin (right) (Yoshida et al., 2009).	18
Fig. 2.17. Pathway for the biosynthesis of phenolics and polyphenols (modified from Salminen and Karonen, 2011).	19
Fig. 2.18. The shikimate pathway (Herrmann and Weaver, 1999).	20
Fig. 2.19. Biosynthesis of phenylalanine and tyrosine (Keller et al., 1982).	21
Fig. 2.20. Biosynthesis of phenolic acids and hydrolyzable tannins (Crozier et al., 2009).....	22
Fig. 2.21. Biosynthesis of flavonoids (Macheix et al., 1990).	23
Fig. 2.22. Biosynthesis of anthocyanidins and flavonols (Bowsher et al., 2008; Gould et al., 2009).	25
Fig. 2.23. Biosynthesis of anthocyanins (Gould et al., 2009; He et al., 2010).	26
Fig. 2.24. Schematic of the probable anthocyanins bioabsorption (McGhie and Stevenson, 2013) and pH values of the different parts in the human gastrointestinal tract (Pandit and Soltis, 2012). Acy: Anthocyanidins; Acy-Gly: anthocyanin glycoside; Acy-methyl: methylated anthocyanidins; Acy-Gluc: anthocyanidin glucuronide.	29
Fig. 2.25. Classification of antioxidants (Carocho and Ferreira, 2013).	33
Fig. 2.26. Diseases induced by oxidative stress in humans (Pham-Huy et al., 2008).	34
Fig. 2.27. Blackberry plant (left, © Wolfgang Dirscherl/pixelio.de) and blackberry fruits (right).....	42
Fig. 2.28. From left to right: Aronia arbutifolia, Aronia prunifolia, Aronia melanocarpa Nero and Viking (Source: with permission of Aronia ORIGINAL Naturprodukte GmbH).....	43
Fig. 2.29. Sweet cherries (© Johannes Heide/pixelio.de) and sour cherries (© Tim Reckmann/pixelio.de).	44



List of figures

Fig. 2.30. Black carrots cultivars: Antonina, Deep Purple, Purple Haze and Purple Sun (From left to right).	45
Fig. 2.31. Purple Sweet Potato from China (left and in the middle) and the distribution of anthocyanin pigmentation in sweet potatoes (right) by Huamán (1992).....	46
Fig. 2.32. Roselle plant from Mexico (left and in the middle, by R. René Rodríguez Arzaba) and its dark red calyces.....	47
Fig. 2.33. Clitoria ternatea L.: seeds (left), flowers (in the middle) and dried flowers (right).	48
Fig. 2.34. Basic model systems for CCC: hydrostatic (left) and hydrodynamic (right) equilibrium systems (Ito and Conway, 1986).	49
Fig. 2.35. Search for the HSCCC (Ito and Conway, 1986).	50
Fig. 2.36. Head (H) to Tail (T) orientations of helical column in synchronous planetary motion. They are determined by the handedness of the coil, right or left (Ito, 1984).....	50
Fig. 2.37. Principle unilateral hydrodynamic equilibrium system (Ito and Conway, 1986).	51
Fig. 2.38. Single-coil design (left) (Ito and Conway, 1986) and HSCCC with multilayer triple-coils (right).....	52
Fig. 2.39. Single-Coil CCC system (modified from Sutherland, 1987).	52
Fig. 2.40. Spectrum HPCCC system which can be used in analytical or semipreparative scale.	53
Fig. 2.41. Low-Speed Rotary Countercurrent Chromatography system (Du et al., 2000).	54
Fig. 2.42. Membrane chromatography system.....	56
Fig. 2.43. Sartobin S IEX 150 mL membrane adsorber capsule (left), microporous structure with pore size 0.45-3µm (at the middle) and flow path (right) (Source: Sartorius AG, Göttingen, Germany).	56
Fig. 3.1. Schema of work.....	57
Fig. 3.2. HPLC-DAD and HPLC-Chromatograms at 520 nm, 280 nm, 320 nm and 360 nm of blackberry. Peak numbering according to Table 3.1, 3.2 and 3.3.	59
Fig. 3.3. Membrane Chromatography overview (a), fraction collection of copigments (b-d) and anthocyanins (e).	63
Fig. 3.4. HPLC-DAD analyses of a separation of a blackberry extract by Membrane Chromatography.	64
Fig. 3.5. HPLC-Chromatograms at 280 nm and 520 nm of copigment fraction and anthocyanin fraction. Peak numbering according (peak 1-4) to Table 3.1 and peak 5 is a cyanidin-derivative.....	65
Fig. 3.6. HPLC-DAD chromatogram of blackberry XAD-7 extract before separation.	66
Fig. 3.7. LSRCCC system: Detector, fraction collector and plotter (left), column and motor (at the middle) and coil with blackberry sample (right).....	66
Fig. 3.8. LSRCCC chromatogram at 520 nm of blackberry XAD-7 extract.	67
Fig. 3.9. Polymeric fraction of blackberry after LSRCCC separation.....	67
Fig. 3.10. DAD-Contour-Plot Chromatogram of LSRCCC Fraction 5 before HSCCC separation (at the top) and HSCCC separation of LSRCCC Fraction 5 at 520 nm (in the middle) and 280 nm (at the bottom).	70
Fig. 3.11. DAD-Contour-Plot Chromatogram and HPLC Chromatograms at 520 nm, 280 nm and 360 nm of Aronia melanocarpa juice. Peak numbering according to Table 3.5.....	72
Fig. 3.12. DAD-Contour-Plot chromatogram and HPLC-chromatograms of Aronia melanocarpa pomace at 520 nm, 280 nm and 374 nm. Peak numbering according to Table 3.5.....	76



List of figures

Fig. 3.13. DAD-Contour-Plot Chromatograms of Aronia melanocarpa pomace XAD-7 extract (at the top), copigment fraction (at the middle) and anthocyanin fraction (at the bottom).	78
Fig. 3.14. HPLC chromatograms of anthocyanin fraction and copigment fraction from Aronia melanocarpa juice extract (A) and pomace (B) at 520 nm and 280 nm. Peak numbering according to Table 3.5.	79
Fig. 3.15. HSCCC chromatogram of the copigment fraction of Aronia melanocarpa pomace at 280 nm.	80
Fig. 3.16. HSCCC chromatogram of the copigment fraction of Aronia melanocarpa juice extract at 280 nm.	81
Fig. 3.17. HSCCC chromatogram of the anthocyanin fraction of Aronia melanocarpa juice extract at 520 nm... ..	83
Fig. 3.18. HSCCC chromatogram of the anthocyanin fraction from Aronia melanocarpa pomace at 520 nm.	83
Fig. 3.19. HPLC chromatogram of anthocyanin fraction from Aronia melanocarpa A20 extract at 520 nm.	84
Fig. 3.20. HPLC-DAD-Contour-Plot chromatogram and Base Peak UV chromatogram at 520 nm of sour cherry extract Actiplants® Cherry Purevital. (For peak numbers see Table 3.8).	85
Fig. 3.21. 5-Carboxypyranocyanidin-3-(2 ^G -glucosylrutinoside) (left) and 5-Carboxypyranocyanidin-3-rutinoside (right).	87
Fig. 3.22. Base Peak UV Chromatogram at 280 nm (at the top), 320 nm (in the middle) and 360 nm (at the bottom) of sour cherry extract. Peak numbering according to Table 3.10.	88
Fig. 3.23. a) Sample loading of Actiplants® Cherry Purevital extract b) elution of anthocyanins and c) elution of copigments after second separation.	91
Fig. 3.24. DAD-Contour-Plot chromatogram of the anthocyanin fraction (a) and of the copigment fraction (b) from Actiplants® Cherry Purevital extract.	91
Fig. 3.25. Base Peak chromatogram at 520 nm and 280 nm of the anthocyanin fraction from sour cherry extract. For peak numbers see Table 3.8.	92
Fig. 3.26. Base Peak chromatogram at 520 nm and 280 nm of the copigment fraction from sour cherry extract. Peak numbering according to Table 3.10.	93
Fig. 3.27. Base Peak Chromatogram at 320 nm of the copigment fraction from sour cherry extract. Peak numbering (Peak 10-19) according to Table 3.10, while peak 20 and peak 21 are 3-FQA and 5-FQA, respectively.	93
Fig. 3.28. DAD-Contour-Plot chromatograms (left) and HPLC chromatograms at 520 nm (right) for detection of anthocyanins of black carrots: a) Antonina, b) Deep Purple, c) Purple Sun, d) Purple Haze and e) Beta Sweet. Peak numbering according to Table 3.14.	95
Fig. 3.29. HPLC chromatograms of black carrots at 280 nm. For peak numbering, cf. Table 3.16.	97
Fig. 3.30. DAD-Contour-Plot chromatograms of the XAD-7 extract, anthocyanin fraction and copigment fraction of Antonina, Deep Purple and Purple Sun carrots.	99
Fig. 3.31. HPLC chromatograms of anthocyanin fraction (A) and copigment fraction (B) from Deep Purple black carrot at 520 nm and 280 nm. Peak numbering according to Table 3.14 and Table 3.16.	99
Fig. 3.32. HPLC chromatograms of anthocyanin fraction (C) and copigment fraction (D) from Antonina black carrot at 520 nm and 280 nm. Peak numbering according to Table 3.14 and Table 3.16.	100
Fig. 3.33. HPLC chromatograms of anthocyanin fraction (E) and copigment fraction (F) from Purple Sun black carrot at 520 nm and 280 nm. Peak numbering according to Table 3.14 and Table 3.16.	100



List of figures

Fig. 3.34. HSCCC chromatograms of anthocyanin fraction from Antonina, Deep Purple and Purple Sun carrots at 520 nm and chemical structure of the main anthocyanin.	102
Fig. 3.35. HSCCC Chromatograms of copigment fraction from Antonina, Deep Purple and Purple Sun carrots at 280 nm.	104
Fig. 3.36. HPLC DAD Plot chromatogram of Chinese PSP (top) and HPLC chromatograms at 520 nm (middle) and 280 nm (bottom). Peak numbering according to Table 3.17 and Table 3.18.	106
Fig. 3.37. Purple Sweet potatoes (a-b), extraction of polyphenols from purple sweet potato (c-d) and isolation of PSP polyphenols by Amberlite XAD-7 column chromatography (e).	109
Fig. 3.38. Separation of purple sweet potato XAD-7 extract by membrane chromatography (left), copigment fraction elution (at the middle) and anthocyanin fraction elution (right).	109
Fig. 3.39. HPLC DAD Plot chromatograms of PSP anthocyanin fraction and PSP copigment fraction.	109
Fig. 3.40. HPLC-DAD chromatograms at 520 nm (A) and 280 nm (B) of copigment and anthocyanin fraction. For peak numbering, see Table 3.17 and Table 3.19.	110
Fig. 3.41. HSCCC chromatogram of PSP anthocyanin fraction at 520 nm.	112
Fig. 3.42. HSCCC chromatogram of PSP copigment fraction at 280 nm.	113
Fig. 3.43. HPLC DAD plot chromatograms of Hibiscus sabdariffa L. from 200 nm to 600 nm and HPL-DAD chromatogram at 520 nm.	114
Fig. 3.44. HPLC-Chromatograms of Hibiscus sabdariffa L. at 280 nm, 320 nm and 360 nm. Peak numbering according to Table 3.21.	116
Fig. 3.45. Hibiscus sabdariffa L.: a) dried calyces, b) extraction of polyphenols, c-d) concentration of polyphenols onto XAD-7 column and e) HPLC DAD plot chromatogram of HSL XAD-7 extract.	118
Fig. 3.46. HPLC DAD plot chromatograms of HSL anthocyanin fraction and HSL copigment fraction after separation by membrane chromatography.	118
Fig. 3.47. HPLC chromatogramms of HSL anthocyanin fraction at 520 nm and 280 nm. For peak numbering, see Table 3.22.	119
Fig. 3.48. HPLC-DAD chromatogramms of HSL copigment fraction at 520 nm, 280 nm, 320 nm and 360 nm. For peak numbering, see Table 3.22.	120
Fig. 3.49. HSCCC separation of Hibiscus sabdariffa L. XAD-7 extract at 520 nm and 280 nm.	123
Fig. 3.50. HPLC DAD plot chromatogram from 200 to 660 nm and Base Peak chromatograms of Clitoria ternatea L. at 520 nm, 360 nm, 320 nm and 280 nm. For peak numbers see Tables 3.23 and 3.24.	125
Fig. 3.51. Biosynthesis of Ternatin C5 (Kogawa et al., 2007).	127
Fig. 3.52. Possible biosynthetic pathways of ternatins and preternatins (Terahara et al., 1998).	127
Fig. 3.53. Clitoria ternatea L.: HPLC-DAD analysis of raw extract (top), anthocyanin fraction (middle) and copigment fraction (bottom).	129
Fig. 3.54. HPLC Chromatogram at 520 nm of raw extract, anthocyanin fraction and copigment fraction of Clitoria ternatea L.	130
Fig. 3.55. HPCCC chromatogram of raw extract of Clitoria ternatea L. at 280 nm.	131
Fig. 3.56. HPCCC chromatogram of copigment fraction of Clitoria ternatea L. at 280 nm.	132
Fig. 3.57. Determination of total phenolic content in red/black fruits.	134



List of figures

Fig. 3.58. HPLC-DAD Profile of *Aronia melanocarpa* juice and aronia extracts AE1-AE4 (left) and detection of adulteration in AE2-AE4 by UV-chromatogram at 520 nm (right)..... 134





List of tables

Table 2.1. Structure of anthocyanidins.	11
Table 2.2. Pharmacokinetics of anthocyanins in humans and animals after consumption (Prior and Wu, 2006; Daayf and Lattanzio, 2009; Fernandes et al., 2015).	30
Table 2.3. Absorption of anthocyanins in cell models (Fernandes et al., 2015).	32
Table 3.1. Anthocyanin content and LC-MS data of anthocyanins in blackberry.	60
Table 3.2. LC-MS data of ellagitannins in blackberry.	61
Table 3.3. Phenolic acids, flavonols and flavan-3-ols in blackberry.	62
Table 3.4. Distribution of phenolic compounds from blackberry after LSRCCC separation.	68
Table 3.5. LC-ESI-MS ⁿ data of polyphenols in Aronia melanocarpa juice.	73
Table 3.6. Anthocyanin content in Aronia melanocarpa products.	75
Table 3.7. Separation of Aronia melanocarpa XAD-7 extracts by Membrane Chromatography.	78
Table 3.8. LC-MS ⁿ data of anthocyanins in sour cherry extract Actiplants® Cherry Purevital.	86
Table 3.9. Anthocyanin content in cherry products.	87
Table 3.10. LC-MS data of copigments in Actiplants® Cherry Purevital extract.	89
Table 3.11. Quantification of chlorogenic acids in Actiplants Cherry Purevital.	90
Table 3.12. Quantification of flavonols in Actiplants Cherry Purevital.	90
Table 3.13. Amount of obtained fractions from sour cherry extract after membrane chromatography.	91
Table 3.14. LC-MS data of the main polyphenols in black carrots.	96
Table 3.15. Quantification of anthocyanins in black carrots.	96
Table 3.16. LC-MS data and quantification of main chlorogenic acids in black carrots.	98
Table 3.17. Mass Spectrometric data and quantification of Anthocyanins from Chinese PSP.	107
Table 3.18. LC-MS data of Copigments in Chinese PSP.	108
Table 3.19. LC-MS data and quantification of phenolic acids present in copigment fraction after separation by membrane chromatography.	111
Table 3.20. LC-MS data and quantification of anthocyanins Hibiscus sabdariffa L. from Mexico.	115
Table 3.21. LC-MS data of phenolic compounds in Hibiscus sabdariffa L.	117
Table 3.22. LC-MS data of phenolic compound in Hibiscus sabdariffa L. after membrane chromatography.	121
Table 3.23. Anthocyanins in Clitoria ternatea L.	126
Table 3.24. Flavonols of Clitoria ternatea L. flowers.	128
Table 4.1. List of raw material.	135
Table 4.2. List of chemicals.	137
Table 4.3. Calibration range for quantification by HPLC.	140
Table 4.4. Technical data of Sartobind SIEX 150 mL, filter capsule and pompe system.	144
Table 4.5. Parameters for CCC separations.	146





Abbreviations

AcOH	Acetic acid
AF	Anthocyanin-fraction
AJ	Concentrated aronia juice
arab	arabinoside
AP	Aronia pomace
A20	Actiplants® Aronia 20 %
Conc.	Concentration
CCC	Counter-current chromatography
CH ₃ CN	Acetonitrile
CF	Copigment-fraction
CQA	Caffeoylquinic acid
Cy	Cyanidin
δ	Chemical shift in ppm (Parts per Million)
d	Doublet
dd	doublet of doublets
ddd	doublet of doublet of doublets
DAD	Diode array detector
DEPT	Distortionless enhancement by polarisation transfer
Del	Delphinidin
diglc	diglucoside
DM	Dry matter
ESI	Electrospray ionization
EtOAc	Ethyl acetate
EtOH	Ethanol
FW	Fresh weight
GAE	Gallic acid equivalent
gal	galactoside
glc	glucoside
H	Head
HCl	Hydrochloric acid
Hz	Hertz
HPLC	High-performance liquid chromatography
HSCCC	High-Speed Counter-current chromatography
i. D.	inside diameter
J	J-coupling in Hertz (s-1; Hz)



Abbreviations

k	Partition coefficient
λ	Wavelength in nm
LC	Liquid Chromatography
LP	Lower phase
LSRCCC	Low Speed Rotary Countercurrent Chromatography
m	Multiplett
M	Molar
[M] ⁺	Pseudo-molecular ion (pos. Modus)
[M-H] ⁻	Pseudo-molecular ion (neg. Modus)
max	Maxima
MC	Membrane chromatography
MeOH	Methanol
min	Minutes
MS	Mass spectrometry
Mv	Malvidin
<i>m/z</i>	Mass-to-charge-ratio
n-BuOH	n-Butanol
n.d.	Not detected
nm	nanometer
NMR	Nuclear magnetic resonance
No.	Number
p. a.	per Analysis
Pg	Pelargonidin
Pn	Peonidin
ppm	Parts per million
Pt	Petunidin
q	quartet, quadruplet
rpm	Revolutions per minute
RP-18	Reversed-phase 18
rut	rutinoside
s	Singlet
samb	sambubioside
soph	sophoroside
T	Tail
t	triplet
tBME	<i>tert</i> -butyl methyl ether
TFA	Trifluoroacetic acid



Abbreviations

TFA-d ₁	Trifluoroacetic acid- d ₁
TLC	Thin-layer chromatography
t _R	Retention time
UV	Ultraviolet
Vis	Visible
v/v	Volume ratio
xyl	xyloside

