




Haruki Iino (Autor)

## Functional analysis of the nuclear basket and protein Tpr



<https://cuvillier.de/de/shop/publications/7587>

Copyright:

Cuvillier Verlag, Inhaberin Annette Jentsch-Cuvillier, Nonnenstieg 8, 37075 Göttingen, Germany  
Telefon: +49 (0)551 54724-0, E-Mail: [info@cuvillier.de](mailto:info@cuvillier.de), Website: <https://cuvillier.de>



# 1. Table of contents

|   |    |
|---|----|
| <b>2. Abstract</b> .....  | 7  |
| <b>3. Introduction</b> .....  | 8  |
| 3.1 Gene expression in eukaryotes   |    |
| 3.2 mRNA processing   |    |
| 3.2.1 Sequential splicing reactions   |    |
| 3.2.2 Alternative splicing  |    |
| 3.2.3 Alternative cleavage and polyadenylation  |    |
| 3.3 mRNA export pathways  |    |
| 3.3.1 NXF1/Tap-mediated export  |    |
| 3.3.2 Crm1-mediated mRNA export   |    |
| 3.4 mRNA quality control (QC)   |    |
| 3.5 Retroviral strategies to bypass mRNA QCs  |    |
| 3.5.1 The RRE-mediated mRNA export  |    |
| 3.5.2 The CTE-mediated export   |    |
| 3.6 The nuclear pore complex  |    |
| 3.7.1 The nuclear basket protein, Tpr   |    |
| 3.7.2 Functional aspects of Tpr   |    |
| 3.8 The aim of this study   |    |
| <b>4. Results</b> .....   | 30 |
| 4.1. Detection of bulk mRNAs upon RNAi of Tpr   |    |
| 4.2. The establishment of a novel reporter system to analyze mRNA export  |    |
| 4.3. The detection of potential un-spliced mRNA leakages in the absence of Tpr using the triple reporter cell lines   |    |
| 4.4. The enrichment of a specific intermediate at the distinct splicing stages does not cause an increase in intron-based reporter activity in the absence of Tpr in interphase cells                         |    |
| 4.5. Studying un-spliced mRNA leakage in the absence of Tpr using an exon-based leakage reporter cell system  |    |
| 4.6. The expression levels of intron-containing HIV- <i>gag</i> mRNAs are elevated by the depletion of Tpr in the presence of the CTE   |    |
| 4.7. Tpr amount required for the HIV- <i>gag</i> mRNA retention   |    |
| 4.8. The effect of exogenously expressed Tpr polypeptides on the HIV-Gag protein levels   |    |
| 4.9. The potential roles of the essential sites required for splicing in the Tpr-dependent HIV- <i>gag</i> mRNA regulation  |    |
| 4.10. The potential role of the HIV- <i>gag</i> ORF sequence in the Tpr-dependent HIV- <i>gag</i> mRNA regulation   |    |
| 4.11. The CTE-containing <i>Photinus luciferase</i> mRNAs are not monitored by Tpr  |    |
| 4.12. Further mapping of the HIV- <i>gag</i> segment involved in the Tpr-dependent nuclear retention of the HIV- <i>gag</i> mRNA  |    |
| 4.13. Investigation on whether Tpr might play a role in the subcellular distribution of other CTE-harboring transcripts, or of some distinct, alternatively spliced, intron-containing endogenous transcripts |    |
| 4.14. Possible explanations for the Tpr-dependent regulation of un-spliced mRNAs  |    |



- 4.14.1. Investigating the potential roles of the HIV inhibitory sequence (INS)-related proteins in the Tpr-dependent HIV-*gag* mRNA retention
- 4.14.2. Investigating a potential contribution of the NPC-binding proteins to the Tpr-dependent HIV-*gag* mRNA regulation
- 4.14.3 RNAi screening: the proteins potentially involved in the HIV-*gag* mRNA regulation
- 4.14.4 Searching for possible contributions of the positive candidates in HIV-*gag* mRNA regulation
- 4.14.5 Investigating a possible contribution of Tpr in the long non-coding RNA degradation pathway
- 4.14.6 Further attempts to gain insight into the Tpr-dependent mechanism of retaining CTE-containing HIV-*gag* transcripts in the nucleus
- 4.15. Proteins potentially involved in the Tpr-dependent mRNA export

## **5. Discussion**..... 91

- 5.1. The establishment of stable reporter cell lines to monitor mRNA retention and export
- 5.2. Considering the possibility of un-spliced mRNAs occurring in mitotic cytosols due to mitotic NE disassembly and the effects on the analysis of mRNA export
- 5.3. A potential contribution of Tpr to bulk mRNA export
- 5.4. Tpr is involved in the regulation of the CTE-containing un-spliced mRNA export
- 5.5 In the Tpr-deficient cell, export of the HIV-*gag*-CTE mRNAs appears to prevent them from being more quantitatively spliced or degraded
- 5.6. Which Tpr pool is responsible for preventing the HIV-*gag*-CTE mRNA export?
- 5.7. How many Tpr molecules are required for the prevention of HIV-*gag*-CTE mRNA export?
- 5.8. How does Tpr prevent the HIV-*gag*-CTE mRNA export?
- 5.9. Which Tpr domain is responsible for the suppression?
- 5.10. The difference of reporter systems between yeast and vertebrates
- 5.11 Perspective

## **6. Materials and Methods**..... 108

## **7. Acknowledgement**..... 123

## **8. Appendix**..... 126

Appendix 1: HIV-*gag* sequences

Appendix 2: Supplement figures

Appendix 3: Table 4-2, and Table 4-3

## **9. References**..... 135