Allan Ndua Mweke (Autor)

Development of entomopathogenic fungi as biopesticides for the management of Cowpea Aphid, Aphis craccivora Koch



https://cuvillier.de/de/shop/publications/7923

Copyright:
Cuvillier Verlag, Inhaberin Annette Jentzsch-Cuvillier, Nonnenstieg 8, 37075 Göttingen, Germany
Telefon: +49 (0)551 54724-0, E-Mail: info@cuvillier.de, Website: https://cuvillier.de

1.0 GENERAL INTRODUCTION

1.1 Background

Cowpea, *Vigna unguiculata* (L.) Walp (Fabaceae) is an important leguminous crop in semi-arid tropics and is well adapted to drought (Akibonde and Maredia, 2011; Horn et al., 2016; Yadav et al., 2017).

Estimated annual production of cowpea is 6.2 million metric tons (MT) occupying 14.5 million hectares of land in more than 45 countries across the world (Abate et al., 2012). Africa is the leading producer of cowpea with Nigeria, Niger and Burkina Faso accounting for 80% of the world's production (FAO, 2016). Kenya produces about 2% of the total world production mainly in the drier regions (HCD, 2014; FAO, 2016).

Consumption of cowpea either as leaves, immature pods, green peas and dry grains provides local populations affordable sources of proteins, essential vitamins and minerals (Ghaly and Alkoaik, 2010; Hall, 2012). Cowpea is used mainly as grain legume in West Africa and as a leafy vegetable in East Africa (Chiulele et al., 2011; Hall, 2012; Rusike et al., 2013). Leafy cowpea is an important African indigenous vegetable (AIV) in Kenya that is widely grown and consumed in urban and rural areas (Abukutsa-Onyango, 2010; Rusike et al., 2013). Cowpea also has medicinal benefits as its consumption has been shown to reduce environmental enteric dysfunction (Trehan et al., 2015).

The vegetative parts of cowpea plant are fed to livestock as fodder (haulms) and farmers earn income from selling them in the dry season (Rusike et al., 2013). The spreading cowpea varieties protect the soil against soil erosion and help in weed control (Garko et al., 2016). The crop improves soil fertility and soil structure by providing organic matter from crop residues and fixing atmospheric nitrogen into the soil through symbiosis with nodule forming bacteria such as *Bradyrhizobium* spp. (Mucheru-Muna, 2010; Schipanski and Drinkwater, 2012; Dwivedi et al., 2015).

1.2 Problem statement and justification of the study

In Africa, yields of cowpea are low, ranging between 100 and 250 kg/ha (Omongo et al., 1997; Baidoo et al., 2012) compared to potential yields of 3000 kg/ha in United States of America (Rusoke and Rubaihayo, 1994; Hall, 2012). Arthropod pests and diseases as well as of use of inferior varieties and/or farmer saved seeds and poor soil fertility are the key limiting factors in cowpea production (Obopile, 2006; Dugje et al., 2009). Cowpea aphid, *Aphis craccivora* Koch (Aphididae), is a major pest of cowpea that attacks the crop in all the stages of its growth (Blackman and Eastop 2006; Kusi et al., 2010; Souleymane et al., 2013). The pest contributes to yield losses of up to 100% through disruption of plant physiological growth

Dieses Werk ist copyrightgeschützt und darf in keiner Form vervielfältigt werden noch an Dritte weitergegeben werden.
Es gilt nur für den persönlichen Gebrauch.

by sacking the plant sap and slowing of photosynthetic process by reducing leaf surface area exposed to light by depositing honeydew on the leaves (Sorensen, 2009). Additional yield losses are also due to transmission of more than 30 plant viruses including Cowpea aphid-borne mosaic virus (CABMV) (Blackman and Eastop, 2000; Smith and Boyko 2007; Damiri et al., 2013).

Management of the cowpea aphid is mainly based on use of chemical pesticides, which have been reported to be expensive and ineffective (Hassan, 2013). Furthermore, pesticides have many undesirable impacts like development of resistance due to indiscriminate use, human health concerns due to exposure during application and chemical residues on plant products, bioaccumulation in the environment and disruption of ecological services of beneficial organisms (Sánchez-Bayo, 2011; Baidoo et al., 2012; El-Heneidy et al., 2015). The negative environmental impacts of the synthetic pesticides have accelerated search for alternative crop protection products leading to increased development of less toxic compounds based on naturally occurring toxins from micro-organisms (Mazid *et al.,* 2011).

Several integrated management approaches targeting cowpea aphid have been developed and used with varying degrees of success (Afun et al., 1991; Egho, 2010). These strategies have employed use of resistant varieties (Huynh et al., 2013; Smith and Chuang, 2014), use of biological control agents including the use of entomopathogenic fungi (EPF) formulated as mycoinsecticides (Hajek and Delalibera, 2010), manipulation of agro ecosystems like intercropping (Hassan, 2013), and targeted insecticide application as opposed to routine and blanket spraying (Egho and Enujeke, 2012). The plant sap sucking soft bodied aphids are susceptible to attack by EPF under natural environment and epizootics due to infection by EPF have been reported (Pell et al., 2003; Roy et al., 2010). The EPF belong to the Zygomycetes and Hyphomycetes groups but the Zygomycetes class form the larger group of EPF attacking aphids including *A. craccivora* (Humber 1991). Several isolates of *Metarhizium anisopliae* (Metschnikoff) Sorokin, *Beauveria bassiana* (Balsamo) Vuillemin, *Lecanicillium* spp and *Isaria* spp are commercial products currently available for use in management of several aphid species. For example, *B. bassiana* (registered as BotaniGard® and Naturalis-L®), *M. anisopliae* (registered as Met52®), *Isaria javanica* (Frieder and Bally), (Samson and Hywel-Jones), (registered as Preferal®) and *Lecanicillium* spp. (registered as Vertalec®) are all used for aphid control in Europe and North America (Zimmerman, 1992; Cook et al., 1996; Whipps, 1997; Fravel et al., 1998; Wraight and Carruthers, 1999; Copping and Menn, 2000; Hynes and Boyetchko, 2006; Jandricic et al., 2014).

Fungal based biopesticides are being used as commercial crop protection products for management of aphids in Asian, Latin American as well as and European countries which account for the greatest market share of these products with Africa registering and using the lowest percentage (Faria and Wraight, 2007). Biopesticides are increasingly becoming viable

Dieses Werk ist copyrightgeschützt und darf in keiner Form vervielfältigt werden noch an Dritte weitergegeben werden.
Es gilt nur für den persönlichen Gebrauch.

alternatives for control of insect pests due to their safety to users, non-target beneficial arthropods and environment and their compatibility with IPM strategies.

Though there are previous studies on pathogenicity of *M. anisopliae*, *B. bassiana* and *Isaria* spp on *A. craccivora* and other aphid species (Ekesi et al., 2000, Sahayaraj and Borgio 2010, Saranya et al., 2010; Bayissa et al., 2016), the studies did not evaluate the promising isolate in this study; ICIPE 62 against *A. craccivora*. Additionally, ICIPE 62 has been used to control other aphid species in vegetables and is commercially available as a biopesticide in Kenya. Moreover, none of the isolates evaluated in this study have been tested for pathogenicity against *A. craccivora* before. In Kenya, there is an EPF based biopesticide (Met 62®- *M. anisopliae* isolate ICIPE 62) developed by the International Centre of Insect Physiology and Ecology (*icipe*) in collaboration with Real IPM Kenya against major vegetable arthropod pests including aphid species (http://www.realipm.com/). However, this product does not include *A. craccivora* as one of the target pests hence this study evaluated the pathogenicity of this isolate among others for potential development of a biopesticide that can be used in the management of this pest.

1.3 General objective

This study aimed at developing and optimizing entomopathogenic fungi as biopesticides for the management of *A. craccivora* within the context of cowpea IPM.

1.4 Specific objectives

1. Screen entomopathogenic fungi isolates for their virulence against the cowpea aphid (*A. craccivora*) and select candidate isolates that can be developed into a biopesticide and used in an IPM system
2. Evaluate different formulations of the selected isolate for the management of *A. craccivora* on cowpea under field conditions
3. Assess the efficacy of intercropping cowpea with maize and application of selected EPF isolate for the management of *A. craccivora*.

1.5 Research questions

This study set out to answer the following questions:

4. Does pathogenicity of entomopathogenic fungi to *Aphis craccivora* vary within and among isolates of different species?
5. Is production of fungal spore on insect cadavers positively related to isolate(s) virulence?
6. Does formulation type influence performance of entomopathogenic isolates both in screenhouse and field conditions?

Dieses Werk ist copyrightgeschützt und darf in keiner Form vervielfältigt werden noch an Dritte weitergegeben werden.
Es gilt nur für den persönlichen Gebrauch.

7. Is combination of intercropping cowpea and maize and application of entomopathogenic fungi more effective in suppressing aphid population under field conditions compared to application of Duduthrin or non-application of either EPF or Duduthrin in a cowpea maize intercrop?

1.6 References

Abate T, Alene AD, Bergvinson D, Shiferaw B, Silim S, Orr A and Asfaw S. 2012. Tropical grain legumes in Africa and South Asia: knowledge and opportunities. International Crops Research Institute for the Semi-Arid Tropics, Nairobi, pp 112.

Abukutsa MOO. 2010. African indigenous vegetables in Kenya: Strategic repositioning in the horticultural sector. Inaugural Lecture, Jomo Kenyatta University of Agriculture and Technology, Nairobi, Kenya. 30 April.

Afun JVK, Jackai LEN and Hodgson CJ. 1991. Calendar and monitored insecticide application for the control of cowpea pests. *Crop Protection*, 10 (5):363-370.

Akibode S and Maredia MK. 2011. Global and regional trends in production, trade and consumption of food legume crops. Department of Agricultural, Food, and Resource Economics Michigan State University. Staff Paper Series, 2012-10.

Baidoo PK, Baidoe-Ansah D and I. Agbonu I. 2012. Effects of Neem (*Azadirachta indica* A. Juss). Products on *Aphis craccivora* and its predator *Harmonia axyridis* on cowpea. *American Journal of Experimental Agriculture*, 2:198-206.

Bayissa W, Ekesi S, Mohamed SA, Kaaya GP, Wagacha JM, Hanna R and Maniania NK. 2016. Selection of fungal isolates for virulence against three aphid pest species of crucifers and okra. *Journal of Pest Science*, 90:355-68.

Blackman RL and Eastop VF. 2000. Aphids on the world's crops: an identification and information guide (No. Ed. 2). John Wiley & Sons Ltd. 476.

Blackman RL and Eastop VF. 2006. Aphids on the world's herbaceous plants and shrubs. Chichester, UK: John Wiley & Sons Ltd, pp. 1460.

Chiulele RM, Mwangi G, Tongoona P, Ehlers JD and Ndeve AD. 2011. Assessment of farmers' perceptions and preferences of cowpea in Mozambique. *African Crop Science Conference Proceedings*, 10:311-318.

Cook RJ, Bruckart WL, Coulson JR, Goettel MS, Humber RA, Lumsden RD, Maddox JV, McManus ML, Moore L, Meyer SF and Quimby Jr PC. 1996. Safety of microorganisms intended for pest and plant disease control: A framework for scientific evaluation, *Biological Control*, 7:333–351.

Copping LG and Menn JJ. 2000. Biopesticides: a review of their action, applications and efficacy. *Pest Management Science*, 56 (8): 651-676.

Damiri BV, Al-Shahwan IM, Al-Saleh MA, Abdalla OA and AmerMA. 2013. Identification and characterization of cowpea aphid-borne mosaic virus isolates in Saudi Arabia. *Journal of Plant Pathology*, 79-85.

Dugje IY, Omoigui LO, Ekeleme F, Kamara AY and Ajeigbe H. 2009. Farmers' guide to cowpea production in West Africa. IITA, Ibadan, Nigeria.

Dieses Werk ist copyrightgeschützt und darf in keiner Form vervielfältigt werden noch an Dritte weitergegeben werden.
Es gilt nur für den persönlichen Gebrauch.

Dwivedi SL, Sahrawat K, Upadhyaya H, Mengoni A, Galardini M, Bazzicalupo M, Biondi EG, Hungria M, Kaschuk G, Blair MW and Ortiz R. 2015. Advances in host plant and rhizobium genomics to enhance symbiotic nitrogen fixation in legumes, in: D.L. Sparks (Ed.), Advances in Agronomy, Elsevier Inc., Amsterdam, pp. 1–116.

Egho EO. 2010. Comparative studies on insect species of cowpea (*Vigna unguiculata*) (L) Walp) in two agro-ecological zones during the early cropping season in Delta State, Southern Nigeria. *Agriculture and Biology Journal of North America*, 1 (5): 946-949.

Egho EO and Enujeke EC. 2012. Minimizing insecticide application in the control of insect pests of cowpea (*Vigna Unguiculata* (L) WALP) in Delta state, Nigeria. *Sustainable Agriculture Research*, 1 (1): 87.

Ekesi S, Akpa AD, Onu I, Ogunlan MO. 2000. Entomopathogenicity of *Beauveria bassiana* and *Metarhizium anisopliae* to the cowpea aphid, *Aphis craccivora*. *Archives of Phytopathology and Plant Protection*, 33:171–180.

El-Heneidy AH, Khidr AA and Taman AA. 2015. Side-effects of insecticides on non-target organisms: 1-In Egyptian Cotton Fields. *Egyptian Journal of Biological Pest Control*, 25 (3):685.

Faria MR and Wraight SP. 2007. Mycoinsecticides and mycoacaricides: a comprehensive list with worldwide coverage and international classification of formulation types. *Biological Control*, 43 (2):237-256.

FAO, 2016. Food and Agriculture Organization of the United Nations, Rome, Italy. http://faostat.fao.org/default.aspx.

Fravel DR, Connick Jr WJ and Lewis JA. 1998. Formulation of microorganisms to control plant diseases. In: Formulation of microbial biopesticides, Springer, Dordrecht, pp. 187-202.

Garko MS, Mohammed IB and Fulani MS. 2016. Performance of cowpea [*Vigna unguiculata* (l.) Walp.] varieties as influenced by weed control treatments in the Sudan Savanna of Nigeria. *International Journal of Scientific and Research Publications*, 6:135-140.

Ghaly AE and Alkoaik FN. 2010. Extraction of protein from common plant leaves for use as human food. *American Journal of Applied Sciences*, 7 (3):331.

Hajek AE and Delalibera I. 2010. Fungal pathogens as classical biological control agents against arthropods. *Bio Control*, 55 (1):147-158.

Hall AE. 2012. Phenotyping cowpeas for adaptation to drought. *Frontiers in Physiology*, 3, 1-8.

Hassan S. 2013. Effect of variety and intercropping on two major cowpea (*Vigna unguiculata* Walp) field pests in Mubi, Adamawa state, Nigeria. *International Journal of Agricultural Research and Development*, 1:108-109.

Horn LN, Ghebrehiwot HM and Shimelis HA. 2016. Selection of novel cowpea genotypes derived through gamma irradiation. *Frontiers in Plant Science*, 7:262.

Horticultural Crops Directorate (HCD). 2014. Annual report.

Dieses Werk ist copyrightgeschützt und darf in keiner Form vervielfältigt werden noch an Dritte weitergegeben werden.
Es gilt nur für den persönlichen Gebrauch.

Humber RA. 1991. Fungal pathogens of aphids. In: Peters DC, Webster JA, Chlouber CS (eds). Aphid-plant interactions: Populations to molecules, 45–56 Agricultural Experiment Station, Division of Agriculture, Oklahoma State University, Stillwater.

Huynh BL, Ehlers JD, Close TJ, Cisse´ N, Drabo I, Boukar O, Lucas MR, Wanamaker S, Pottorff M and Roberts PA. 2013. Enabling tools for modern breeding of cowpea for biotic stress resistance Translational genomics for crop breeding. Wiley, London, pp. 183–199.

Hynes RK and Boyetchko SM. 2006. Research initiatives in the art and science of biopesticide formulations. *Soil Biology and Biochemistry*, 38 (4):845-849.

Jandricic SE, Filotas M, Sanderson JP and Wraight SP. 2014. Pathogenicity of conidia-based preparations of entomopathogenic fungi against the greenhouse pest aphids *Myzus persicae*, *Aphis gossypii*, and *Aulacorthum solani* (Hemiptera: Aphididae). *Journal of Invertebrate Pathology*, 118: .34-46.

Kusi F, Obeng-Ofori D, Asante SK and Padi FK. 2010. New sources of resistance in cowpea to the cowpea aphid (*Aphis craccivora* Koch) (Homoptera: Aphididae). *Journal of the Ghana Science Association*, 12 (2): 95-104.

Mazid S, Kalida JC and Rajkhowa RC. 2011. A review on the use of biopesticides in insect pest management. *International Journal of Science and Advanced Technology*, 1 (7):169-178.

Mucheru-Muna M, Pypers P, Mugendi D, Kung'u J, Mugwe J, Merckx R and Vanlauwe B. 2010. A staggered maize–legume intercrop arrangement robustly increases crop yields and economic returns in the highlands of Central Kenya. *Field Crops Research*, 115 (2):132-139.

Obopile M. 2006. Economic threshold and injury levels for control of cowpea aphid, *Aphis craccivora* Linnaeus (Homoptera: Aphididae) on cowpea. *African Plant Protection*, 12:111–115.

Omongo CA, Ogenga-Latigo MW, Kyamanywa S and Adipala E. 1997. The effect of seasons and cropping systems on the occurrence of cowpea pests in Uganda. In *African Crop Science Conference Proceedings*, 3:1111-1116.

Pell JK, Eilenberg J, Hajek AE and Steinkraus DC. 2001. Biology, ecology and pest management potential of Entomophthorales. In: Butt TM, Jackson C, Magan N (eds) Fungi as biocontrol agents: progress, problems and potential. CAB International, Wallingford, pp. 71–153.

Roy HE, Brodie EL, Chandler D, Goettel MS, Pell JK, Wajnberg E and Vega F. 2010. Deep space and hidden depths: understanding the evolution and ecology of fungal entomopathogens. *Biological Control*, 55:1–6.

Rusike JG, van den Brand S, Boahen K, Dashiell S, Kantengwa J, Ongoma DM, Mongane G, Kasongo ZB, Jamagani R, Aidoo R and Abaidoo R. 2013. Value chain analyses of grain legumes in Kenya, Rwanda, Eastern DRC, Ghana, Nigeria, Mozambique, Malawi and Zimbabwe. (No. 1.2. 6, 1.3. 4). N2Africa.

Dieses Werk ist copyrightgeschützt und darf in keiner Form vervielfältigt werden noch an Dritte weitergegeben werden.
Es gilt nur für den persönlichen Gebrauch.

Rusoke DG and Rubaihayo PR. 1994. The influence of some crop protection management practices on yield stability of cowpea. *African Crop Science Journal*, 2:43-48.

Sahayaraj K and Borgio JF. 2010. Virulence of entomopathogenic fungus *Metarhizium anisopliae* (Metsch.) Sorokin on seven insect pests. *Indian Journal of. Agricultural Science*, 44:195–200.

Sánchez-Bayo F. 2011. Impacts of agricultural pesticides on terrestrial ecosystems. In: Sánchez-Bayo, F, van den Brink, PJ, Mann, R (eds.) Ecological Impacts of Toxic Chemicals. Bentham Science Publishers, Online, pp. 63-87.

Saranya SR, Ushakumari S, Jacob S and Philip BM. 2010. Efficacy of different entomopathogenic fungi against cowpea aphid, *Aphis craccivora* (Koch). *Journal of Biopesticides*, 3:1138–142.

Schipanski ME and Drinkwater LE. 2012. Nitrogen fixation in annual and perennial legume-grass mixtures across a fertility gradient. *Plant and Soil*, 357 (1-2):147-159.

Smith CM and Boyko EV. 2007. The molecular bases of plant resistance and defense responses to aphid feeding: current status. *Entomologia Experimentalis et Applicata*, 122 (1):1-16.

Smith CM and Chuang WP. 2014. Plant resistance to aphid feeding: behavioral, physiological, genetic and molecular cues regulate aphid host selection and feeding. *Pest Management Science*, 70:528–540.

Souleymane A, Aken'Ova ME, Fatokun CA and Alabi OY. 2013. Screening for resistance to cowpea aphid (*Aphis craccivora* Koch) in wild and cultivated cowpea (*Vigna unguiculata* Walp.) accessions. *International Journal of Science, Environment and Technology*, 2 (4): 611 – 621.

Trehan I, Benzoni NS, Wang AZ, Bollinger LB, Ngoma TN, Chimimba UK, Stephenson KB, Agapova SE, Maleta KM and Manary MJ. 2015. Common beans and cowpeas as complementary foods to reduce environmental enteric dysfunction and stunting in Malawian children: study protocol for two randomized controlled trials. *Trials*, 16 (1):520.

Whipps JM. 1997. Developments in the biological control of soil-borne plant pathogens. *Advances in Botanical Research*, 26:1–134.

Wraight SP and Carruthers RL. 1999. Production, delivery, and use of mycoinsecticides for control of insect pests of field crops. In: Hall FR, Menn JJ, (eds). Methods in Biotechnology, Vol. 5: Biopesticides: Use and Delivery. Totowa, New Jersey, USA: Humana Press, pp. 233-69.

Yadav T, Nisha KC, Chopra NK, Yadav MR, Kumar R, Rathore DK, Soni PG, Makarana G, Tamta A, Kushwah M and Ram H. 2017. Weed management in cowpea-A Review. *International Journal of Current Microbiology and Applied Sciences* 6 (2):1373-1385.

Zimmermann G. 2007 Review on safety of the entomopathogenic fungus *Metarhizium anisopliae. Biocontrol Science and Technology*, 17 (9):879-920.

Dieses Werk ist copyrightgeschützt und darf in keiner Form vervielfältigt werden noch an Dritte weitergegeben werden.
Es gilt nur für den persönlichen Gebrauch.

2.0 SCIENTIFIC BACKGROUND

2.1 *Vigna unguiculata* L. Walp

Origin: Cowpea (*Vigna unguiculata* L. Walp.) is an annual drought tolerant legume that is adapted to different soil types and different cropping systems whose origin has been traced to Southern Africa region (Singh et al., 1997). The West African countries including Nigeria, Niger, Burkina Faso, Benin, Togo and Cameroon have the biggest diversity of cultivated cowpea (Ng and Marechal; 1985; Ng, 1995; Padulosi and Ng, 1997; Timko and Singh, 2008).

Global production*:* Cowpea (*Vigna unguiculata* L. Walp).is grown mostly in tropical Africa (Ofuya, 1997) and Africa produces the largest proportion of the world's production with Nigeria, Niger and Burkina Faso producing 80% of world's production (Mortimore et al., 1997; FAO, 2016). In 2016, the world production was 6,991,174 metric tons with a total value of U$ 18 billion while the total area under production was 12.3 million hectares. In the same year Africa produced 6,739,689 metric tons representing 96% of the world production (FAO, 2016).

Production in Kenya*:* Major cowpea production localities in Kenya are largely in the arid and semi-arid areas since it is a drought tolerant crop and farmers can harvest even when cereal crops like maize and sorghum fail (Saidi et al. 2010). In 2016, the total area under production was 31,020 hectares producing 115,801tons valued at Kenyan shillings 2.4 billion representing about 2% of the world production with Makueni County producing 35% of the total production (HCD, 2016). Areas of production include Machakos, Kitui, Makueni, Tharaka, Mbeere, Taita Taveta, Kwale, Kilifi, Lamu, Kisii, Migori, Homabay, Siaya, Kisumu and Bugoma The cowpea varieties grown by farmers can be categorize into 4 according to their seed colour or mode of growth (1) the cream types which have cream colored seeds, (2) the crowder type with black spots or brown colour, (3) the black eye types whose seeds are white with black eye and (4) other types with intermediate colours. Some commercial varieties in Kenya and East African region include Kunde M66, Ken Kunde 1(KK1) Ken Kunde 3(KK3), K80, KVU 27-31 and KVU 419.

Uses*:* Cowpea crop has a wide adaptability to different climatic conditions and it is cultivated in warm regions of the world mainly for its edible seed, however, the crop is also an important source of vegetable and, it is one of the most important African leafy vegetables (Hall, 2012; Rusike et al., 2013). Cowpea does well in poor soils because it has tolerance to low soil fertility and it has ability to fix atmospheric nitrogen in association with root nodule forming bacteria (*Bradyrhizobium* spp (Schipanski and Drinkwater LE 2012; Ddamulira et al., 2015) and it also able tolerate a wide range of soil pH (Mucheru-Muna, 2010).

In Kenya and other East African countries, cowpea has wide nutritional and agronomic uses. Young leaves are used as a vegetable while the seeds and young pods constitute a rich source

Dieses Werk ist copyrightgeschützt und darf in keiner Form vervielfältigt werden noch an Dritte weitergegeben werden.
Es gilt nur für den persönlichen Gebrauch.

of dietary protein. Dried shoots and roots are used as fuel (Hall, 2012; Trehan et al., 2015). Cowpea seed is a nutritious component in the human diet as it contains about 25% protein and 64% carbohydrate, and 2% fat (Akibode, 2011; Owolabi et al., 2012).The leaves have higher protein content compared to seed (Baker et al. 1989; Nielsen et al., 1993). Cowpea is widely consumed all over the world, mainly in rural populations, and satisfy a considerable proportion of the protein requirements (Oiye et al., 2009; Ghaly and Alkoaik, 2010).

Above ground parts of cowpea plant excluding the pods are used as fodder for livestock (haulms) and a source of income for farmers who harvest and sell during dry period (Singh et al., 1997). The spreading and indeterminate or semi-determinate varieties of cowpea provide ground cover against soil erosion while at the same time suppressing weeds (Singh et al., 1997; Mucheru-Muna, 2010). The cowpea crop residues when ploughed into the soil provide organic matter that improves soil fertility (Mucheru-Muna, 2010). Cowpea being a leguminous crop fixes atmospheric nitrogen into the soil through symbiosis with nodule forming bacteria (*Bradyrhizobium* spp), (Singh et al., 1997; Schipanski and Drinkwater, 2012; Dwivedi et al., 2015).

2.2 The Cowpea aphid *Aphis craccivora* (Koch)

Taxonomic description*:* Compared to other aphid species, *A. craccivora* is a relatively small aphid. Apterous females are characterized by black or dark brown body, brown to yellow legs and a prominent cauda. Nymphs are waxy compared to adults. Adults are distinguished from other closely related aphid species by the presence of 6-segmented antennae with black distal part of femur, siphunculi and cauda (Blackman and Eastop, 2006). Winged (alate) females of *A. craccivora* have distinct dorsal cross bars on the abdomen (Blackman and Eastop, 2000).

Distribution and occurrence: *Aphis craccivora* is widely distributed in the world and it has been reported in regions/countries where it was absent mainly due to changing climate but it is more endemic in the tropics (Blackman and Eastop, 2000).

Biology and ecology

Aphis craccivora has a wide distribution in the tropics where females reproduce parthenogenetically but sexual morphs have been reported in temperate regions (Blackman and Eastop, 2007). Cowpea aphid females are ovoviviparous and retain their eggs inside their bodies and give birth to nymphs. Small colonies of *A. craccivora* establish on actively growing plant parts like leaves, tips and young stems and are frequently found in association with ants (Flatt and Weisser, 2000; Espadaler et al., 2012). A number of biotypes of *A. craccivora* have identified (Ofuya, 1997; Sorensen, 2009).

Development of *A. craccivora* is influenced by climatic conditions including temperature (24-28.5°C), relative humidity 65% RH, hours of sunshine (day length-L: D 16:8) and

Dieses Werk ist copyrightgeschützt und darf in keiner Form vervielfältigt werden noch an Dritte weitergegeben werden.
Es gilt nur für den persönlichen Gebrauch.

precipitation (Mayeux, 1984). Host plant biochemistry like low levels of hydrocarbon positively influences development of alate individuals (Mayeux, 1984). The lifespan of adult's ranges between 5-15 days and under favourable weather conditions *A. craccivora* completes a generation in 10 to 20 days. An adult aphid can produce 20 nymphs in one day and developmental period between first instar and adult is between 3-5 days while a single adult female can produce up to 100 nymphs in their life time (Ofuya, 1997). Weather conditions, soil moisture content and fertility as well as host plant status influence growth, development, reproduction and the lifespan of *A. craccivora* (Ofuya 1997).

Host range

Aphis craccivora is a highly polyphagous aphid species feeding on Leguminoseae group of plants including cowpea (*Vigna unguicalata* (L.) (Walp.), groundnut (*Arachis hypogaea* L.), mungbean (*Vigna radiata* (L.) R. Wilczek), pigeonpea (*Cajanus cajan* (L.) Huth), chickpea (*Cicer arietinum* L.), green beans (*Vicia* spp. and *Phaseolus* spp.), lupins (*Lupinus angustifolius* L.), lentil (*Lens esculenta*) and lucerne (*Medicago sativa* L.). It is also reported as a minor pest on other leguminous non-leguminous crops, such as cotton and citrus (Blackman and Eastop, 2006; Brady and White, 2013).

Economic importance of *Aphis craccivora*

Direct crop damage: *Aphis craccivora* causes direct damage by sucking plant sap and injecting toxins into the phloem during all plant growth stages including seedlings, flowers and pods and this damage is by both adults and nymphs (Ofuya, 1997; Huynh et al., 2015). When heavy infestations occur during early plant growth stages young plants wither and eventually die and those that survive are characterized by stunted growth, distorted leaves and experience delayed flowering and lower yields (Ofuya, 1995). Heavy infestation by cowpea aphid at podding stage can reduce seed yield (Ofuya, 1997).

Indirect damage*:* High population of *A. craccivora* produces high amounts of honeydew on plant leaf surfaces thereby promoting growth of the sooty mold fungus which interferes with respiratory and photosynthesis capacity of the plant by altering biochemical and physiological processes of infested plants thereby reducing plant growth and associated yield (Gomez et al., 2004; Sorensen, 2009; Goławska et al., 2010). Honeydew also reduces quality of cowpea leaves and renders them inedible thus contributing to yield loss. *Aphis craccivora* is a known vector of more 30 plant viruses including cowpea mosaic virus, ground nut rosette virus (GRV), subterranean clover stunt virus (SCSV) (clover stunt virus), Bean common mosaic virus (BCMV) (bean mosaic virus, bean western mosaic virus, mungbean mosaic virus) (Atiri et al., 1986; Blackman and Eastop, 2000; Brault et al., 2010).

Dieses Werk ist copyrightgeschützt und darf in keiner Form vervielfältigt werden noch an Dritte weitergegeben werden.
Es gilt nur für den persönlichen Gebrauch.