Contents

Contents I

List of manuscripts III

List of relevant abbreviations IV

Chapter 1: Prologue 1

1.1 Legumes in cropping systems 1
1.2 Intercropping systems 1
1.2.1 Intercropping with faba bean 2
1.2.2 Intercropping with white clover and chicory 3
1.3 Drought and water use 4
1.3.1 Water use efficiency on the physiological level 4
1.3.2 Water use efficiency of the crop stand via remote sensing 5
1.4 Objectives and structure of the thesis 6

Chapter 2 13

Chapter 2.1: The cropping system matters – Contrasting responses of winter faba bean (*Vicia faba* L.) genotypes to drought stress 14

Chapter 2.2: Crop genotype and plant compartment determine the response of the active bacterial community towards water deficit 44

Chapter 2.3: Response of the active bacterial and fungal communities in the rhizosphere differ towards water deficit 83

Chapter 2.4: Crop species and cropping system alter the effect of *Metarhizium brunneum* seed application on plant-associated bacterial and fungal communities 119

Chapter 3: Intercropping with winter faba bean in arable land and white clover in grassland improves water use efficiency of the cropping system 158

Chapter 4: Legume-based mixed intercropping systems may lower agricultural born N_2O emissions 184

Chapter 5: Remotely assessed vegetation development and water use of arable and grassland intercropping with legumes 194

Chapter 6: Epilogue 223

6.1 Intercropping with arable crops 223
6.2 Intercropping with grassland crops 224
6.3 Drought stress tolerance 225
6.4 Breeding for intercropping 226
6.4.1 Genotypes of winter faba bean 227
6.4.2 Genotypes of white clover 228
6.5 Technical aspects 229
6.6 Concluding remarks 230

Summary 235

Further publications 238

Acknowledgments 240

Appendix 242

Supplementary material for chapter 2.1 242
Supplementary material for chapter 2.2 259
Supplementary material for chapter 2.3 260
Supplementary material for chapter 2.4 261
Supplementary material for chapter 3 266

CV 274

Declarations 276