



Elfadl Yousif Elmogtaba Elfadl (Autor)

Suitability of Safflower (*Carthamus tinctorius* L.) for Cultivation as an Oil Crop under the Conditions of Organic and Low-Input Farming Systems in Central Europe

Elfadl Yousif Elmogtaba Elfadl

Suitability of Safflower (*Carthamus tinctorius* L.) for Cultivation as an Oil Crop under the Conditions of Organic and Low-Input Farming Systems in Central Europe



Cuvillier Verlag Göttingen
Internationaler wissenschaftlicher Fachverlag

<https://cuvillier.de/de/shop/publications/836>

Copyright:

Cuvillier Verlag, Inhaberin Annette Jentsch-Cuvillier, Nonnenstieg 8, 37075 Göttingen, Germany

Telefon: +49 (0)551 54724-0, E-Mail: info@cuvillier.de, Website: <https://cuvillier.de>

Table of Contents

Table of contents

Table of contents.....	i
List of Tables.....	iii
List of Figures	v
List of abbreviations.....	vi
1 General introduction	1
1.1 Safflower origins and history	2
1.2 Safflower production and economy	3
1.3 Safflower genomics.....	4
1.4 Safflower as a multi-purpose crop	4
1.5 General safflower characteristics	6
1.6 Production requirements	6
1.7 Germplasm diversity	8
1.8 Breeding for organic safflower	9
1.8.1 Breeding for high oil yield	10
1.8.2 Diseases resistance	11
1.8.3 Weed control	11
1.8.4 Winter hardiness.....	12
1.8.5 Seed dormancy	12
1.8.6 Safflower hybrid development.....	13
1.9 Objectives.....	13
1.10 References	14
2 Evaluation of phenotypic variation in a worldwide germplasm collection of safflower (<i>Carthamus tinctorius</i> L.) grown under organic farming conditions in Germany	23
2.1 Introduction	24
2.2 Materials and Methods.....	25
2.3 Results	28
2.4 Discussion	35
2.5 Conclusions	41
2.6 References	42
3 Safflower (<i>Carthamus tinctorius</i> L.) as an alternative oil crop in organic farming system: Potential and stability in Central Europe.....	47
3.1 Introduction	48
3.2 Materials and Methods	50

Table of Contents

3.3	Results and Discussion.....	54
3.4	Conclusion.....	66
3.5	References	66
4	Optimization of nitrogen rate and seed density for safflower (<i>Carthamus tinctorius</i> L.) production under low-input farming conditions in temperate climate	73
4.1	Introduction	73
4.2	Materials and methods	75
4.3	Results	78
4.4	Discussion	86
4.5	Conclusions	98
4.6	References	98
5	Development of near infrared reflectance spectroscopy (NIRS) calibration model for estimation of oil content in a worldwide safflower germplasm collection	105
5.1	Introduction	106
5.2	Materials and Methods	107
5.3	Results and Discussion.....	109
5.4	Conclusions	116
5.5	Acknowledgements	116
5.6	References	117
6	General discussion and conclusions.....	121
6.1	Pattern of variation	121
6.2	Multivariate analysis	125
6.3	Yield stability	127
6.4	Optimization of nitrogen rate and seed density.....	128
6.5	Development of NIRS calibration for oil estimation	131
6.6	Ideotyping spring safflower for organic farming.....	132
6.7	Conclusions	136
7	Summary	137
8	Zusammenfassung.....	140
9	General references.....	145
10	Appendix.....	169
11	Acknowledgements.....	171
12	Curriculum Vitae.....	173

Table of Contents

List of Tables

Table 2.1: Variance components, overall means, coefficients of variations, and heritabilities for phenotypic traits and rated diseases in safflower germplasm.....	30
Table 2.2: Eigenvector, Eigenvalue, total variance and cumulative variance for 200 accessions and 11 geographic regions based on 10 phenotypic traits and 3 diseases in safflower.....	32
Table 2.3: Phenotypic and genotypic coefficients of correlation (r) between different traits for 468 safflower accessions grown under organic farming conditions during the growing seasons of 2004 and 2005 in South West Germany.....	39
Table 3.1: Variance components, ranges, means, genotypic coefficient of variation (GCV) and heritability (HER) of 20 genotypes (G) tested across 8 environments (E) in the growing seasons of 2004 and 2005	55
Table 3.2: Eigenvector, eigenvalue, total variance and cumulative variance for twenty genotypes based on 14 phenotypic traits and 3 rated diseases in safflower.....	56
Table 3.3: Analysis of variance of yield (t/ha) and oil yield (kg/ha) for 20 genotypes tested over 13 environments in Central Europe across 2 years (2004, 2005)	60
Table 3.4: Means, stability estimates of yield (dt/ha) and ranks of 20 genotypes tested across 13 environments in Central Europe in seasons of 2004 and 2005	60
Table 3.5: Means, standard errors, stability estimates of oil yield (kg/ha) and overall ranks of 20 genotypes evaluated across 13 environments in Central Europe in seasons of 2004 and 2005	62
Table 3.6: Spearman's rank correlation coefficients between stability models for seed yield (below diagonal) and oil yield (above diagonal) of 20 safflower genotypes evaluated over 13 environments in seasons of 2004 and 2005	63
Table 3.7: Phenotypic (above diagonal) and genotypic (below diagonal) coefficients of correlation between different traits for 20 safflower genotypes grown under organic farming conditions averaged over five environments during the growing seasons of 2004 and 2005.....	65
Table 4.1: Long term and growing seasons (2004 and 2005) monthly mean temperatures (°C) and total of precipitation (mm) at Ihinger Hof (Iho) and Wil	76
Table 4.2: Pre-fertilization and post-harvest soil nitrogen levels (NO ₃ -N kg/ha) in 0-90 cm depth at Ihinger Hof (Iho) during 2004 and 2005 growing seasons.....	79
Table 4.3: Factorial variance components of environment (E), nitrogen rate (N), seed density (D), cultivar (C), and their interactions for morpho-phenological traits and rated	

Table of Contents

diseases and agronomic traits data collected at Ihinger Hof (2004 and 2005) and Wil (2004)	84
Table 4.4: Effects and mean comparisons (LSD) of nitrogen rate (N), seed density (D) and cultivar (C) on assessed morpho-phenological traits and rated diseases at Ihinger Hof (Iho) and Wil during 2004 and 2005 growing seasons	90
Table 4.5: Effects and means comparisons (LSD) of environment (E), nitrogen rate (N), seed density (D) and cultivar (C) on oil yield (kg/ha) and its components, biomass yield (kg/ha), and harvest index at Ihinger Hof (Iho) and Wil during 2004 and 2005 growing seasons	94
Table 4.6: Correlation coefficients of oil yield, biomass yield and selected oil yield components based on adjusted means of three factors (N, D, C) applied across 2004 and 2005 growing seasons at Ihinger Hof (Iho)	97
Table 5.1: Calibration and validation statistics in NIRS models for estimation of oil content (%) in ground seed of safflower for combined and individual years	111