Table of contents

Table of contents i
List of Tables iii
List of Figures v
List of abbreviations vi
1 General introduction 1
1.1 Safflower origins and history 2
1.2 Safflower production and economy 3
1.3 Safflower genomics 4
1.4 Safflower as a multi-purpose crop 4
1.5 General safflower characteristics 6
1.6 Production requirements 6
1.7 Germplasm diversity 8
1.8 Breeding for organic safflower 9
1.8.1 Breeding for high oil yield 10
1.8.2 Diseases resistance 11
1.8.3 Weed control 11
1.8.4 Winter hardiness 12
1.8.5 Seed dormancy 12
1.8.6 Safflower hybrid development 13
1.9 Objectives 13
1.10 References 14
2 Evaluation of phenotypic variation in a worldwide germplasm collection of safflower (*Carthamus tinctorius* L.) grown under organic farming conditions in Germany 23
2.1 Introduction 24
2.2 Materials and Methods 25
2.3 Results 28
2.4 Discussion 35
2.5 Conclusions 41
2.6 References 42
3 Safflower (*Carthamus tinctorius* L.) as an alternative oil crop in organic farming system: Potential and stability in Central Europe 47
3.1 Introduction 48
3.2 Materials and Methods 50

3.3 Results and Discussion 54
3.4 Conclusion 66
3.5 References 66
4 Optimization of nitrogen rate and seed density for safflower (*Carthamus tinctorius* L.) production under low-input farming conditions in temperate climate 73
4.1 Introduction 73
4.2 Materials and methods 75
4.3 Results 78
4.4 Discussion 86
4.5 Conclusions 98
4.6 References 98
5 Development of near infrared reflectance spectroscopy (NIRS) calibration model for estimation of oil content in a worldwide safflower germplasm collection 105
5.1 Introduction 106
5.2 Materials and Methods 107
5.3 Results and Discussion 109
5.4 Conclusions 116
5.5 Acknowledgements 116
5.6 References 117
6 General discussion and conclusions 121
6.1 Pattern of variation 121
6.2 Multivariate analysis 125
6.3 Yield stability 127
6.4 Optimization of nitrogen rate and seed density 128
6.5 Development of NIRS calibration for oil estimation 131
6.6 Ideotyping spring safflower for organic farming 132
6.7 Conclusions 136
7 Summary 137
8 Zusammenfassung 140
9 General references 145
10 Appendix 169
11 Acknowledgements 171
12 Curriculum Vitae 173

List of Tables

Table 2.1: Variance components, overall means, coefficients of variations, and heritabilities for phenotypic traits and rated diseases in safflower germplasm..... 30

Table 2.2: Eigenvector, Eigenvalue, total variance and cumulative variance for 200 accessions and 11 geographic regions based on 10 phenotypic traits and 3 diseases in safflower..... 32

Table 2.3: Phenotypic and genotypic coefficients of correlation (r) between different traits for 468 safflower accessions grown under organic farming conditions during the growing seasons of 2004 and 2005 in South West Germany..... 39

Table 3.1: Variance components, ranges, means, genotypic coefficient of variation (GCV) and heritability (HER) of 20 genotypes (G) tested across 8 environments (E) in the growing seasons of 2004 and 2005 55

Table 3.2: Eigenvector, eigenvalue, total variance and cumulative variance for twenty genotypes based on 14 phenotypic traits and 3 rated diseases in safflower..... 56

Table 3.3: Analysis of variance of yield (t/ha) and oil yield (kg/ha) for 20 genotypes tested over 13 environments in Central Europe across 2 years (2004, 2005) 60

Table 3.4: Means, stability estimates of yield (dt/ha) and ranks of 20 genotypes tested across 13 environments in Central Europe in seasons of 2004 and 2005 60

Table 3.5: Means, standard errors, stability estimates of oil yield (kg/ha) and overall ranks of 20 genotypes evaluated across 13 environments in Central Europe in seasons of 2004 and 2005 62

Table 3.6: Spearman's rank correlation coefficients between stability models for seed yield (below diagonal) and oil yield (above diagonal) of 20 safflower genotypes evaluated over 13 environments in seasons of 2004 and 2005 63

Table 3.7: Phenotypic (above diagonal) and genotypic (below diagonal) coefficients of correlation between different traits for 20 safflower genotypes grown under organic farming conditions averaged over five environments during the growing seasons of 2004 and 2005..... 65

Table 4.1: Long term and growing seasons (2004 and 2005) monthly mean temperatures (°C) and total of precipitation (mm) at Ihinger Hof (Iho) and Wil 76

Table 4.2: Pre-fertilization and post-harvest soil nitrogen levels (NO3-N kg/ha) in 0-90 cm depth at Ihinger Hof (Iho) during 2004 and 2005 growing seasons..... 79

Table 4.3: Factorial variance components of environment (E), nitrogen rate (N), seed density (D), cultivar (C), and their interactions for morpho-phenological traits and rated

diseases and agronomic traits data collected at Ihinger Hof (2004 and 2005) and Wil (2004) 84

Table 4.4: Effects and mean comparisons (LSD) of nitrogen rate (N), seed density (D) and cultivar (C) on assessed morpho-phenological traits and rated diseases at Ihinger Hof (Iho) and Wil during 2004 and 2005 growing seasons 90

Table 4.5: Effects and means comparisons (LSD) of environment (E), nitrogen rate (N), seed density (D) and cultivar (C) on oil yield (kg/ha) and its components, biomass yield (kg/ha), and harvest index at Ihinger Hof (Iho) and Wil during 2004 and 2005 growing seasons 94

Table 4.6: Correlation coefficients of oil yield, biomass yield and selected oil yield components based on adjusted means of three factors (N, D, C) applied across 2004 and 2005 growing seasons at Ihinger Hof (Iho) 97

Table 5.1: Calibration and validation statistics in NIRS models for estimation of oil content (%) in ground seed of safflower for combined and individual years 111