

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 3 |
| 2 | The antiferromagnetic $S = 1/2$ Heisenberg chain model | 8 |
| 2.1 | Hamiltonian | 8 |
| 2.2 | Thermodynamic properties of the isotropic AFHC model | 14 |
| 2.2.1 | Magnetization | 14 |
| 2.2.2 | Specific heat | 16 |
| 2.3 | Dynamical structure factor | 17 |
| 2.3.1 | Field and temperature dependence of $S(q, \omega)$ | 18 |
| 2.4 | Determination of $1/T_1$ by means of conformal field theory | 24 |
| 3 | Nuclear magnetic resonance | 31 |
| 3.1 | Magnetization and susceptibility | 31 |
| 3.2 | Static hyperfine fields | 34 |
| 3.2.1 | Types of hyperfine fields | 35 |
| 3.2.2 | Dipole-dipole coupling | 36 |
| 3.2.3 | Fermi contact exchange | 38 |
| 3.2.4 | Shielding tensor | 38 |
| 3.2.5 | Quadrupole interaction | 39 |
| 3.2.6 | Rotation of magnetic hyperfine fields | 41 |
| 3.3 | Level splitting in the presence of hyperfine fields | 41 |
| 3.3.1 | The nuclear Zeeman effect - magnetic splitting | 41 |
| 3.3.2 | Quadrupolar splitting | 43 |
| 3.4 | Dynamic hyperfine fields | 44 |
| 3.4.1 | Magnetic fluctuations - The Bloch equations | 45 |
| 3.4.2 | Measurement of relaxation times - pulse sequences | 47 |
| 3.4.3 | Measurement of T_2 - spin decoherence | 48 |
| 3.4.4 | Measurement of T_1 - scattering with quasiparticles | 49 |
| 3.5 | Relaxation in the presence of Zeeman and quadrupolar splitting | 50 |
| 3.6 | Spin-lattice relaxation caused by magnetic fluctuations | 52 |
| 3.7 | Spin-spin relaxation due to indirect coupling | 53 |
| 3.8 | Instrumental setups | 53 |

| | |
|--|------------|
| 4 The spin chain systems CuPzN and PhnCuCl | 57 |
| 4.1 Copper Pyrazine Dinitrate (CuPzN) | 57 |
| 4.1.1 Magnetic exchange mechanisms | 60 |
| 4.1.2 Sample preparation | 62 |
| 4.1.3 Review of experiments from the literature | 62 |
| 4.2 Bisphenazinium Copper Tetrachloride (PhnCuCl) | 71 |
| 4.2.1 Crystal structure | 71 |
| 4.2.2 Sample preparation | 73 |
| 5 Results and discussion of the thermodynamic experiments | 76 |
| 5.1 Copper Pyrazine Dinitrate | 76 |
| 5.1.1 Magnetization | 76 |
| 5.2 Bisphenazinium Copper Tetrachloride | 78 |
| 5.2.1 Magnetization | 78 |
| 5.2.2 Specific heat | 85 |
| 5.2.3 Conclusion | 88 |
| 6 Results and discussion of the NMR experiments | 90 |
| 6.1 Copper Pyrazine Dinitrate | 90 |
| 6.1.1 ^{13}C NMR: Static properties and hyperfine coupling mechanisms | 92 |
| 6.1.2 ^{13}C NMR: Comparison of dynamics with QMC calculations | 103 |
| 6.1.3 ^{13}C NMR: Comparison of dynamics with analytical theory | 110 |
| 6.1.4 ^{14}N NMR: Static properties and hyperfine coupling | 114 |
| 6.1.5 Conclusion | 124 |
| 6.2 Bisphenazinium Copper Tetrachloride | 126 |
| 6.2.1 Calibration of the cryomagnet | 127 |
| 6.2.2 ^1H NMR: Static properties and hyperfine coupling | 130 |
| 6.2.3 ^1H NMR: Field- and temperature-dependent dynamics | 137 |
| 6.2.4 ^{35}Cl NMR: Comparison with the ^1H results | 141 |
| 6.2.5 Discussion of four typical first-approach scenarios | 144 |
| 6.2.6 Conclusion | 145 |
| 7 Summary | 147 |