

Table of contents

1. General Introduction	1
1.1 References	3
2. Foraging behaviour of steers grazing a heterogeneous sward - Effects of breed and stocking density	5
2.1 Abstract	5
2.2 Introduction	6
2.3 Materials and methods	7
2.3.1 Treatments and management	8
2.3.2 Behaviour Observation	9
2.3.3 Bite and Step Rate	10
2.3.4 Daylight Quarters	10
2.3.5 Statistical analysis	10
2.4 Results	12
2.4.1 Behaviour patterns	12
2.4.2 bpm, spm and bps	14
2.5 Discussion	15
2.5.1 Daylight quarter	15
2.5.2 Period	16
2.5.3 Year	17
2.5.4 Breed	18
2.5.5 Stocking density	21
2.6 Conclusions	23
2.7 References	24
3. Selectivity of grazing on heterogeneous grassland - The effect of cattle breed and stocking density	29
3.1 Abstract	29
3.2 Introduction	30
3.3 Material and methods	31
3.3.1 Treatments and management	31
3.3.2 Bite types	33
3.3.3 Bite types on offer	33
3.3.4 Consumption of bite types	34
3.3.5 Quality of herbage on offer	34
3.3.6 Jacobs' Index	35
3.3.7 Statistical analysis	35
3.4 Results	37
3.4.1 Bite types on offer	37
3.4.2 Quality of herbage on offer	39
3.4.3 Selectivity	42
3.5 Discussion	44
3.5.1 Period	45
3.5.2 Daylight quarter	46
3.5.3 Stocking density	48
3.5.4 Breed	49
3.6 Conclusions	49
3.7 References	51

4. Performance of beef cattle steers at continuous grazing on grassland managed for biodiversity benefits	57
4.1 Abstract	57
4.2 Introduction	58
4.3 Materials and methods.....	59
4.3.1 Treatments and management	60
4.3.2 CSH and dry matter on offer.....	61
4.3.3 Animal weight.....	61
4.3.4 Pasture Performance.....	62
4.3.5 Statistical analysis.....	63
4.4 Results.....	64
4.4.1 Compressed sward height	64
4.4.2 Management and performance	65
4.5 Discussion	69
4.5.1 Stocking	69
4.5.2 Breed	72
4.5.3 Suggestions for developing maintenance and production systems	73
4.6 Conclusions	75
4.7 References.....	77
5. General Discussion.....	81
5.1 References.....	85
6. Summary	87
7. Zusammenfassung	89
8. Acknowledgements	93