

<u>CONTENTS</u>	Page
1 INTRODUCTION.....	1
2 MATERIAL AND METHODS.....	5
2.1 Laboratory experiments.....	5
2.1.1 Rearing of the insects.....	5
2.1.1.1 Rearing of <i>Frankliniella occidentalis</i> and <i>Thrips tabaci</i> .....	5
2.1.1.1.1 Stock culture.....	5
2.1.1.1.2 Obtaining of individuals in the desired stage.....	5
2.1.1.2 Rearing of collembolan and natural enemies.....	6
2.1.2 Cultivation entomopathogenic fungi.....	7
2.1.3 Procedures of the experiments.....	7
2.1.3.1 Screening efficiency isolates of different entomopathogenic fungi against thrips species, <i>Frankliniella occidentalis</i> and <i>Thrips tabaci</i> .....	7
2.1.3.1.1 Preliminary screening according to pathogenicity.....	9
2.1.3.1.2 Secondary screening according to virulence degree.....	9
2.1.3.1.3 Tertiary screening according to effectiveness.....	10
2.1.3.2 Biological and ecological characteristics of the selected high efficiency <i>Beauveria bassiana</i> Bb.5335 and <i>Metarhizium anisopliae</i> Ma.7965....	10
2.1.3.2.1 Morphological characteristics.....	10
2.1.3.2.2 Physiological characteristics.....	12
2.1.3.2.2.1 Evaluation of conidia viability.....	12
2.1.3.2.2.2 Germination at different environmental factors	12
2.1.3.2.2.2.1 Temperature.....	12
2.1.3.2.2.2.2 Relative humidity.....	13
2.1.3.2.2.2.3 Light-dark conditions.....	13
2.1.3.2.2.2.4 Ultraviolet light.....	14
2.1.3.2.2.3 Conidiation at different environmental factors	14
2.1.3.2.2.3.1 Temperature.....	14
2.1.3.2.2.3.2 Relative humidity.....	15
2.1.3.2.2.3.3 Light-dark conditions.....	15
2.1.3.2.2.3.4 Ultraviolet light.....	16

---

2.1.3.2.2.4	Colony growth at different environmental factors.....	16
2.1.3.2.2.4.1	Temperature.....	16
2.1.3.2.2.4.2	Relative humidity.....	17
2.1.3.2.2.4.3	Light-dark conditions.....	17
2.1.3.2.2.4.4	Ultraviolet light.....	17
2.1.3.3	Mode of infection of <i>Beauveria bassiana</i> Bb.5335 and <i>Metarhizium anisopliae</i> Ma.7965 to <i>Frankliniella occidentalis</i> .....	18
2.1.3.3.1	Attachment of conidia.....	18
2.1.3.3.2	Germination and evaluation of germ tube on cuticle.....	19
2.1.3.3.3	Penetration of fungal into the cuticle .....	19
2.1.3.3.4	Colonization.....	20
2.1.3.4	Efficiency of <i>Beauveria bassiana</i> Bb.5335 and <i>Metarhizium anisopliae</i> Ma.7965 against <i>Frankliniella occidentalis</i> and <i>Thrips tabaci</i> .....	20
2.1.3.4.1	Influence of abiotic factors.....	20
2.1.3.4.2	Influence of host plants.....	21
2.1.3.5	Side effects on non-target arthropods.....	22
2.2	Experiments under greenhouse conditions.....	25
2.2.1	Persistence of <i>Beauveria bassiana</i> Bb.5335 and <i>Metarhizium anisopliae</i> Ma.7965 conidia on different host plants.....	25
2.2.1.1	Conidia viability after prolonged exposure.....	26
2.2.1.2	Efficiency of conidia after prolonged exposure.....	26
2.2.2	Efficiency of <i>Beauveria bassiana</i> Bb.5335 and <i>Metarhizium anisopliae</i> Ma.7965 for controlling <i>Frankliniella occidentalis</i> in greenhouse.....	27
2.3	Statistical analysis.....	27
3	RESULTS.....	29
3.1	Laboratory experiments.....	29
3.1.1	Screening efficiency isolates of different entomopathogenic fungi against thrips species, <i>Frankliniella occidentalis</i> and <i>Thrips tabaci</i> .....	29
3.1.1.1	Preliminary screening according to pathogenicity.....	29
3.1.1.2	Secondary screening according to virulence degree.....	31
3.1.1.3	Tertiary screening according to effectiveness.....	33

3.1.2	Biological and ecological characteristics of the selected high efficiency <i>Beauveria bassiana</i> Bb.5335 and <i>Metarhizium anisopliae</i> Ma.7965.....	36
3.1.2.1	Morphological characteristics.....	36
3.1.2.2	Physiological characteristics.....	38
3.1.2.2.1	Evaluation of conidia viability.....	38
3.1.2.2.2	Germination at different environmental factors.....	39
3.1.2.2.2.1	Temperature.....	39
3.1.2.2.2.2	Relative humidity.....	40
3.1.2.2.2.3	Light-dark conditions.....	41
3.1.2.2.2.4	Ultraviolet light.....	42
3.1.2.2.3	Conidiation at different environmental factors.....	43
3.1.2.2.3.1	Temperature.....	43
3.1.2.2.3.2	Relative humidity.....	44
3.1.2.2.3.3	Light-dark conditions.....	45
3.1.2.2.3.4	Ultraviolet light.....	45
3.1.2.2.4	Colony growth at different environmental factors.....	46
3.1.2.2.4.1	Temperature.....	46
3.1.2.2.4.2	Relative humidity.....	47
3.1.2.2.4.3	Light-dark conditions.....	47
3.1.2.2.4.4	Ultraviolet light.....	48
3.1.3	Mode of infection of <i>Beauveria bassiana</i> Bb.5335 and <i>Metarhizium anisopliae</i> Ma.7965 to <i>Frankliniella occidentalis</i> .....	49
3.1.3.1	Attachment of conidia .....	49
3.1.3.2	Germination and evaluation of germ tube on cuticle.....	51
3.1.3.3	Penetration of fungal into the cuticle.....	55
3.1.3.4	Colonization.....	59
3.1.4	Efficiency of <i>Beauveria bassiana</i> Bb.5335 and <i>Metarhizium anisopliae</i> Ma.7965 against <i>Frankliniella occidentalis</i> and <i>Thrips tabaci</i> .....	61
3.1.4.1	Influence of abiotic factors.....	61
3.1.4.1.1	Temperature.....	61
3.1.4.1.2	Relative humidity.....	63
3.1.4.2	Influence of host plants.....	65

---

3.1.5	Side effects on non-target arthropods.....	66
3.2	Experiments under greenhouse conditions.....	68
3.2.1	Persistence of <i>Beauveria bassiana</i> Bb.5335 and <i>Metarhizium anisopliae</i> Ma.7965 conidia on different host plants.....	68
3.2.1.1	Conidia viability after prolonged exposure .....	68
3.2.1.1.1	Crop plants.....	68
3.2.1.1.2	Ornamental plants.....	68
3.2.1.2	Efficiency of conidia after prolonged exposure.....	69
3.2.1.2.1	Crop plants.....	69
3.2.1.2.2	Ornamental plants.....	71
3.2.1.2.3	Lethal time (LT <sub>50</sub> ).....	72
3.2.2	Efficiency of <i>Beauveria bassiana</i> Bb.5335 and <i>Metarhizium anisopliae</i> Ma.7965 for controlling <i>Frankliniella occidentalis</i> in greenhouse.....	73
4	DISCUSSION.....	75
	SUMMARY.....	93
	REFERENCES.....	97