




Shimin Tian (Autor)

Development and application of enzymatic and immunological methods for differentiation and quantification of *Septoria tritici* and *Stagonospora nodorum* in winter wheat and winter triticale



<https://cuvillier.de/de/shop/publications/2803>

Copyright:

Cuvillier Verlag, Inhaberin Annette Jentsch-Cuvillier, Nonnenstieg 8, 37075 Göttingen, Germany

Telefon: +49 (0)551 54724-0, E-Mail: info@cuvillier.de, Website: <https://cuvillier.de>

	CONTENTS	I
	ABBREVIATIONS	IV
1	INTRODUCTION	1
2	MATERIALS AND METHODS	8
2.1	Chemicals	8
2.2	Culture media	9
2.3	Fungal species and isolates	10
2.4	Fungal culture and protein extraction	11
2.5	Experiments in greenhouse	12
2.5.1	Cultivation of plant	12
2.5.2	Inoculums and inoculation	12
2.5.3	Sampling and preparation	12
2.5.4	Visual assessment	13
2.6	Experiment under field conditions	13
2.6.1	Fungicide experiments	13
2.6.2	Variety experiments	14
2.7	Enzymatic methods	14
2.7.1	Extraction buffer	14
2.7.2	Enzyme preparation in liquid culture	14
2.7.3	Enzyme preparation from plant materials	15
2.7.4	Measurement of enzyme activity	15
2.7.4.1	Plate - clearing assay	16
2.7.4.2	Quantitative enzyme assay	16
2.8	Determination of protein concentration	17
2.9	Immunological methods	18
2.9.1	Antigen extraction from plant materials	18
2.9.2	Fungal protein preparation	19
2.9.3	Immunoantigen preparation	19
2.9.4	Immunisation of rabbits	20
2.9.5	Purification of antiserum	21
2.9.6	Biotinylation of antibody	21
2.9.7	Procedures of ELISA	22

2.10	Statistical analysis	25
3	RESULTS	27
3.1	Enzymatic methods for detection of <i>S. tritici</i> and <i>S. nodorum</i>	27
3.1.1	Enzyme production of <i>S. tritici</i> and <i>S. nodorum</i>	27
3.1.1.1	On agar diffusion test	27
3.1.1.2	Enzyme production in liquid culture	28
3.1.1.3	Enzyme activities in infected wheat leaves	30
3.1.2	Relationship between enzyme activity and disease severity	32
3.1.2.1	Enzyme activity in infected leaves and disease severity under greenhouse conditions	32
3.1.2.2	Xylanase and cellulase activities and <i>Septoria</i> leaf diseases under field conditions	33
3.1.3	Relationship between enzyme activity and fungicide efficacy and yield	34
3.1.4	Enzyme activity in different wheat varieties under natural infection conditions	38
3.2	Immunological methods for detection of <i>S. tritici</i> and <i>S. nodorum</i>	41
3.2.1	Selection of <i>S. tritici</i> antisera	41
3.2.1.1	Comparison of the antisera by indirect ELISA	41
3.2.1.2	Comparison of the antibodies in BA-ELISA	42
3.2.2	Reactivity of St-IgG954 and Sn-IgG958	44
3.2.2.1	Optimisation of the antibodies concentration	44
3.2.2.2	Standardisation and sensitivity of the antibodies	45
3.2.2.3	Specificity of the antibodies with fungal proteins	47
3.2.2.4	Reactivity of the antibodies with plant extracts	49
3.2.3	Qualitative detection of <i>S. tritici</i> and <i>S. nodorum</i>	50
3.2.3.1	Early detection of the fungi in wheat leaves in greenhouse tests	50
3.2.3.2	Early detection of the fungi in wheat leaves under field conditions	50
3.2.4	Quantification of <i>S. tritici</i> and <i>S. nodorum</i>	52
3.2.4.1	Extraction methods for fungal antigens from plants	52
3.2.4.2	Quantification of the fungi in wheat leaves from greenhouse tests	52
3.2.4.3	Quantification of the fungi in wheat leaves under field conditions	54

3.2.5	Fungicide effects on <i>S. tritici</i> and <i>S. nodorum</i>	56
3.2.5.1	Early assessments of fungicide efficacy	56
3.2.5.2	Progression of the pathogens in wheat leaves	58
3.2.5.3	Fungicide efficacy and the relationship to yield	60
3.2.6	Variety resistance to <i>S. tritici</i> and <i>S. nodorum</i>	64
3.2.6.1	Detection of the fungal infections in different wheat tissues	64
3.2.6.2	Resistance of wheat and triticale varieties under field conditions	64
4	DISCUSSION	70
5	SUMMARY	88
6	REFERENCES	91
7	APPENDIX	100