




Xuenong Xu (Autor)

**Combined releases of predators for biological control of spider mites *Tetranychus urticae* Koch and western flower thrips *Frankliniella occidentalis* (Pergande)**



<https://cuvillier.de/de/shop/publications/2870>

Copyright:  
Cuvillier Verlag, Inhaberin Annette Jentsch-Cuvillier, Nonnenstieg 8, 37075 Göttingen,  
Germany  
Telefon: +49 (0)551 54724-0, E-Mail: [info@cuvillier.de](mailto:info@cuvillier.de), Website: <https://cuvillier.de>

## Contents

<b>1.</b>	<b>Introduction.....</b>	<b>17</b>
<b>2.</b>	<b>Interactions in the biological control of western flower thrips <i>Frankliniella occidentalis</i> (Pergande) and two-spotted spider mite <i>Tetranychus urticae</i> Koch with the predatory mite <i>Amblyseius cucumeris</i> Oudemans on beans.....</b>	<b>23</b>
2.1	Introduction.....	23
2.2	Materials and methods.....	24
2.2.1	Plant material.....	24
2.2.2	Insects and mites.....	24
2.2.3	Experimental procedures.....	25
2.2.4	Statistical analysis.....	26
2.3	Results.....	28
2.3.1	Experiment I.....	28
2.3.1.1	Effects of <i>A. cucumeris</i> on TSSM.....	28
2.3.1.2	Effects of <i>A. cucumeris</i> on WFT.....	28
2.3.1.3	Effects of <i>A. cucumeris</i> on TSSM in the presence of WFT.....	29
2.3.1.4	Effects of <i>A. cucumeris</i> on WFT in the presence of TSSM.....	30
2.3.2	Experiment II: Predation of WFT larvae on <i>A. cucumeris</i> eggs.....	32
2.4	Discussion.....	32
<b>3.</b>	<b>Interactions in the biological control of western flower thrips <i>Frankliniella occidentalis</i> (Pergande) and two-spotted spider mite <i>Tetranychus urticae</i> Koch by the predatory bug <i>Orius insidiosus</i> Say on beans.....</b>	<b>37</b>
3.1	Introduction.....	37
3.2	Materials and methods.....	38
3.2.1	Plant material.....	38
3.2.2	Insects and mites.....	38
3.2.3	Experimental procedures.....	39
3.2.4	Data analyses.....	40
3.3	Results.....	41
3.3.1	Effects of <i>O. insidiosus</i> on the TSSM population.....	41

3.3.2	Effects of <i>O. insidiosus</i> on WFT .....	42
3.3.3	Effects of <i>O. insidiosus</i> on the TSSM population in the presence of WFT.....	42
3.3.4	Effects of <i>O. insidiosus</i> on WFT in the presence of TSSM.....	44
3.3.5	Influence of WFT and TSSM on the reproduction of <i>O. insidiosus</i> .....	46
3.4	Discussion.....	47
<b>4.</b>	<b>The effects of combined releases of the predatory mite <i>Amblyseius cucumeris</i> Oudemans and the predatory bug <i>Orius insidiosus</i> Say on mixed infestations of western flower thrips <i>Frankliniella occidentalis</i> (Pergande) and two-spotted spider mite <i>Tetranychus urticae</i> Koch .....</b>	<b>51</b>
4.1	Introduction.....	51
4.2	Materials and methods.....	53
4.2.1	Plant material.....	53
4.2.2	Insects and mites.....	53
4.2.3	Experimental procedures.....	54
4.2.4	Data analyses.....	55
4.3	Results.....	56
4.3.1	Effects of <i>A. cucumeris</i> and <i>O. insidiosus</i> on TSSM populations in the presence of WFT .....	56
4.3.1.1	TSSM population suppression by both predators .....	56
4.3.1.2	Effects of the predator and pest densities on the percentages of TSSM suppression.....	57
4.3.1.3	Proportion of TSSM active stages (larvae, nymphs) and eggs to the total offspring of TSSM under the influence of both predators.....	59
4.3.1.4	Effects of the predator and prey densities on the proportions of TSSM active stages (larvae, nymphs) and eggs.....	60
4.3.1.5	Total number of TSSM adults and the proportion of live TSSM adults on plants at the end of the experiment.....	61
4.3.1.6	Effect of the predator and prey densities on the proportion of live TSSM adults on plants at the end of the experiment.....	63
4.3.2	Effects of <i>A. cucumeris</i> and <i>O. insidiosus</i> on the corrected	

mortality of WFT in the presence of TSSM.....	64
4.3.2.1 Effects of both predators on the corrected mortality of WFT in the presence of TSSM.....	64
4.3.2.2 Effects of the densities of <i>A. cucumeris</i> and <i>O. insidiosus</i> on the corrected mortality of WFT in the presence of TSSM.....	65
4.3.3 Fluctuation of <i>A. cucumeris</i> population.....	66
4.3.3.1 Effects of both predators and pests on the number of offspring per <i>A. cucumeris</i> female.....	66
4.3.3.2 Total number of adults and proportion of live <i>A. cucumeris</i> adults on the plant at the end of the experiment.....	69
4.3.4 The daily number of offspring per <i>O. insidiosus</i> female.....	72
4.4 Discussion.....	72
<b>5. Biocontrol of western flower thrips <i>Franklinilla occidentalis</i> (Pergande) by combined releases of plant-inhabiting predatory mite, <i>Amblyseius cucumeris</i> Oudemans or bug, <i>Orius insidiosus</i> Say with soil-dwelling mite, <i>Hypoaspis aculeifer</i> Canestrini in the presence of two-spotted spider mite <i>Tetranychus urticae</i> Koch.....</b>	<b>77</b>
5.1 Introduction.....	77
5.2 Materials and methods.....	79
5.2.1 Plant material.....	79
5.2.2 Insects and mites.....	79
5.2.3 Experimental procedures.....	79
5.2.4 Data analysis.....	80
5.3. Results.....	81
5.3.1 Experiment 1.....	81
5.3.1.1 Effects of <i>A. cucumeris</i> on TSSM.....	81
5.3.1.2 Thrips control by <i>A. cucumeris</i> and <i>H. aculeifer</i> alone and in combination.....	82
5.3.2 Experiment 2.....	82
5.3.2.1 Effects of <i>O. insidiosus</i> on the TSSM population.....	82
5.3.2.2 Thrips control by <i>O. insidiosus</i> and <i>H. aculeifer</i> alone and in combination.....	83

5.4.	Discussion.....	84
6.	<b>General discussion.....</b>	<b>87</b>
7.	<b>References.....</b>	<b>93</b>
8.	<b>Acknowledgements.....</b>	<b>110</b>