



Patcharin Krutmuang (Autor)

**Detection of genes differentially expressed in  
porcine leucocytes due to transport stress by using  
cDNA-AFLP and Differential Display**



Arbeiten aus dem  
Institut für Tierzuchtwissenschaft  
Rheinische Friedrich-Wilhelms-Universität zu Bonn

Patcharin Krutmuang

Detection of genes differentially expressed  
in porcine leucocytes due to transport stress  
by cDNA-AFLP and Differential Display

Heft: 121

<https://cuvillier.de/de/shop/publications/3101>

Copyright:

Cuvillier Verlag, Inhaberin Annette Jentzsch-Cuvillier, Nonnenstieg 8, 37075 Göttingen,  
Germany

Telefon: +49 (0)551 54724-0, E-Mail: [info@cuvillier.de](mailto:info@cuvillier.de), Website: <https://cuvillier.de>

Contents	Page
Abstract	III
List of abbreviations	IX
List of tables	XIII
List of figures	XIV
1 Introduction	1
2 Literature review	3
2.1 Stress	3
2.1.1 Transportation stress in pigs	3
2.1.2 Stress response mechanisms in animals	5
2.2 Stress indicators in pigs	6
2.3 Genetics and stress response in pigs	8
2.4 Molecular genetic approaches to investigate stress response genes in pigs	9
2.4.1 Identification of the stress response genes	9
2.4.1.1 Quantitative trait loci analysis	10
2.4.1.2 Differential display	10
2.4.1.3 cDNA-AFLP	11
2.4.2 Stress gene localization	13
2.4.2.1 Physical mapping	13
2.4.2.2 Genetic mapping	14
2.4.2.3 Comparative mapping	15
2.4.3 Quantification of gene expression	15
2.4.4 Real time quantitative PCR technique	16
3 Materials and methods	17
3.1 Materials	17

3.1.1	Animals	17
3.1.2	Chemicals, biological materials, reagents and medium	17
3.1.2.1	Chemicals	17
3.1.2.2	Biological materials	19
3.1.2.3	Reagents and media	20
3.1.3	Used software	22
3.1.4	Equipments	22
3.2	Methods	23
3.2.1	Experimental design, blood sampling and RNA isolation	24
3.2.1.1	Experimental design	24
3.2.1.2	Blood sampling	24
3.2.1.3	RNA isolation	25
3.2.2	Detection of differential expressed genes	25
3.2.2.1	Pooled RNA samples	25
3.2.2.2	cDNA-AFLP	27
3.2.2.3	Differential display reverse transcription polymerase chain reaction (DDRT-PCR)	30
3.2.2.4	Characterization the nucleotide sequence of the differentially expressed bands	32
3.2.3	Mapping	36
3.2.3.1	Somatic cell hybrid mapping	36
3.2.3.2	Porcine radiation hybrid mapping	37
3.2.4	Validation of differential bands	39
3.3	Statistical analysis	40
4	Results	42
4.1	Identification of differentially expressed ESTs in leucocytes by cDNA-AFLP	42
4.1.1	Expression profile of ESTs from cDNA-AFLP	42
4.1.2	Homologies to known genes and clone identification	43

4.2	Identification of differentially expressed ESTs in leucocytes by DDRT-PCR	46
4.2.1	Expression profile and characterization of the cloned bands from DDRT-PCR	47
4.3	Mapping of candidate genes expressed during transport stress	52
4.3.1	Regional assignment using somatic cell hybrid panel	52
4.3.2	Radiation hybrid panel	52
4.3.3	Comparative mapping	54
4.4	Validation of cDNA-AFLP and differential display by real time PCR	57
4.4.1	Specificity and linearity of the real time PCR detection	60
4.4.2	Relative quantification of cDNA-AFLP	62
4.4.3	Real time RT-PCR assay validation of differential display results	67
5	Discussion	70
5.1	Expression profiling due to transport stress	71
5.1.1	Expression profile of ESTs stress based on day of transport	72
5.1.2	Expression profile of ESTs stress based on cortisol response levels	73
5.2	Clone identification	74
5.2.1	Homologies to known genes in GenBank	75
5.2.2	Validation of differential gene expression by quantitative real time PCR	76
5.2.3	Function of genes	78
5.2.3.1	Function of candidate genes related to stress response (agreement with quantitative real time PCR results)	78
5.2.3.2	Function of potential candidate gene related to stress response based on sequence information (without validation by real time PCR)	82
5.3	Mapping	84
5.3.1	Confirmation the result of IMpRH map by comparative mapping of human and pigs	85

5.4	Future prospects for investigation of candidate genes for transport stress in pigs	88
6	Summary	89
7	Zusammenfassung	92
8	References	95