




Stefan Hohnwald (Autor)

A Grass-Capoeira Pasture Fits Better Than a Grass-Legume Pasture in the Traditionale Agricultural System of Smallholdings in the Brazilian Humid Tropics



<https://cuvillier.de/de/shop/publications/3358>

Copyright:

Cuvillier Verlag, Inhaberin Annette Jentsch-Cuvillier, Nonnenstieg 8, 37075 Göttingen, Germany

Telefon: +49 (0)551 54724-0, E-Mail: info@cuvillier.de, Website: <https://cuvillier.de>

TABLE OF CONTENTS

	Page
LIST OF TABLES.....	V
LIST OF FIGURES.....	VI
ABBREVIATIONS.....	VIII
1. INTRODUCTION.....	1
2. LITERATURE REVIEW.....	3
2.1. The Capoeira within the slash-and-burn cycle.....	3
2.2. Pastures in the Brazilian humid tropics.....	8
2.3. Ways to prevent ecological degradation of pastures.....	13
2.3.1. Grass-Legume pasture.....	14
2.3.2. Grass-Capoeira pasture.....	22
2.4. Development of hypothesis.....	29
3. MATERIALS AND METHODS.....	31
3.1. Study location.....	31
3.2. Study site and plot design.....	33
3.3. Pasture establishment and management.....	35
3.4. Data collection and analysis.....	41
3.4.1. Indicators for area restoration.....	41
3.4.2. Below ground restoration.....	42
3.4.3. Above ground restoration.....	46
3.4.4. Cattle performance.....	48
4. RESULTS.....	49
4.1. Below ground restoration.....	49
4.1.1. Soil fertility.....	49
4.1.2. Soil bulk-density.....	63
4.2. Above ground restoration.....	63
4.2.1. Phytodiversity.....	64
4.2.2. Soil cover percentage.....	67
4.2.3. Similarity.....	69

4.2.4. Direct cattle impact.....	72
4.2.5. Representativeness of the Capoeira.....	78
4.3. Cattle performance.....	81
5. DISCUSSION.....	83
6. SUMMARY.....	99
7. RESUMO (PORTUGUESE).....	102
8. ZUSAMMENFASSUNG (GERMAN).....	105
9. REFERENCES.....	108
10. APPENDIX.....	122
11. ACKNOWLEDGEMENTS.....	135