

Table of contents

| | | |
|-----------|--|-----------|
| 1. | Introduction | 1 |
| 1.1. | Mitochondria and energy metabolism | 1 |
| 1.2. | Cytochrome c oxidase | 1 |
| 1.2.1. | Structure of cytochrome c oxidase | 1 |
| 1.2.2. | Catalytic activity of cytochrome c oxidase | 4 |
| 1.2.3. | Nucleotide binding sites of cytochrome c oxidase | 6 |
| 1.2.4. | Allosteric ATP-inhibition of cytochrome c oxidase | 6 |
| 1.2.5. | Reversible cAMP-dependent phosphorylation of cytochrome c oxidase | 7 |
| 1.2.6. | Variable H ⁺ /e ⁻ stoichiometries in cytochrome c oxidase | 8 |
| 1.2.7. | High ATP/ADP-ratios decrease the H ⁺ /e ⁻ stoichiometry of only heart cytochrome c oxidase | 10 |
| 1.3. | Regulation of cell respiration | 10 |
| 1.3.1. | The first mechanism of respiratory control | 10 |
| 1.3.2. | The second mechanism of respiratory control | 11 |
| 1.4. | Aims of the present dissertation | 13 |
| 2. | Materials | 14 |
| 2.1. | Tissues | 14 |
| 2.2. | Bacteria and vector | 14 |
| 2.3. | Primers | 14 |
| 2.4. | Apparatuses | 14 |
| 2.5. | Kits | 15 |
| 2.6. | Chromatographic materials | 15 |
| 2.7. | Membranes | 15 |
| 2.8. | Chemicals | 15 |
| 2.9. | Lipids, fatty acids, and derivatives | 15 |
| 2.10. | Detergents | 16 |
| 2.11. | Dyes | 16 |
| 2.12. | Antibodies | 16 |
| 2.13. | Nucleotides | 16 |
| 2.14. | Substrates | 16 |
| 2.15. | Uncouplers and reductants | 16 |
| 2.16. | Enzymes | 17 |
| 2.17. | Antibiotic and inhibitor | 17 |
| 3. | Methods | 18 |
| 3.1. | Molecular biological methods | 18 |
| 3.1.1. | General methods | 18 |
| | Preparation of LB medium | 18 |
| | Agarose gel electrophoresis of DNA | 19 |
| | DNA precipitation by Na-acetate and ethanol | 19 |
| | Photometric quantitation of DNA and RNA | 20 |
| 3.1.2. | RNA preparation | 20 |
| | General instruction for RNA preparation | 20 |
| | Purification of total RNA | 20 |
| 3.1.3. | Complementary DNA (cDNA) preparation | 21 |
| | Reverse transcription (RT) | 21 |

| | | |
|---------|---|----|
| | Purification of cDNA after RT-PCR | 22 |
| 3.1.4. | Polymerase Chain Reaction (PCR) | 23 |
| | Standard-PCR | 23 |
| | Hot-Start-PCR | 24 |
| | Touch-Down-PCR | 24 |
| | 3'-rapid amplification of cDNA ends-PCR (3'-RACE-PCR) | 24 |
| 3.1.5. | Preparation of DNA for cloning | 26 |
| | Elution of DNA from agarose gel | 26 |
| | Refilling of dsDNA-ends with Klenow DNA polymerase | 26 |
| | Phosphorylation of DNA | 27 |
| 3.1.6. | Preparation of vector plasmid for cloning | 27 |
| | Dephosphorylation with shrimp alkaline phosphatase | 29 |
| 3.1.7. | Preparation of competent cells for electrotransformation | 29 |
| 3.1.8. | Ligation of DNA with plasmid | 29 |
| 3.1.9. | Transfer of plasmid DNA into <i>E. coli</i> by electroporation | 30 |
| 3.1.10. | Rapid preparation of plasmid DNA by the TENS method | 30 |
| 3.1.11. | Digestion of plasmid DNA with Pvu II | 31 |
| 3.1.12. | DNA sequencing | 31 |
| 3.2. | Protein chemical methods | 32 |
| 3.2.1. | Isolation of mitochondria | 33 |
| 3.2.2. | Purification of cytochrome c oxidase by the Triton X-100 method | 34 |
| | Extraction by Triton X-114 and Triton X-100 | 35 |
| | DEAE-Sephacel anion exchange chromatography | 35 |
| | Purification of cytochrome c oxidase by fractionated ammonium sulfate precipitation | 35 |
| 3.2.3. | Isolation of cytochrome c oxidase by the cholate method | 36 |
| | Isolation of Keilin-Hartree particles | 36 |
| | Isolation of cytochrome c oxidase | 36 |
| 3.2.4. | Spectrophotometric determination of cytochrome concentrations | 39 |
| | Difference spectrum | 39 |
| | Absolute spectrum | 39 |
| 3.2.5. | Determination of protein concentration | 40 |
| 3.2.6. | SDS-polyacrylamide gel electrophoresis (SDS-PAGE) | 41 |
| | Staining of protein bands with Coomassie Blue | 42 |
| 3.2.7. | Labelling of SH-groups in cytochrome c oxidase with DACM | 42 |
| 3.2.8. | Electroblotting of proteins onto nitrocellulose and PVDF membrane | 43 |
| | Staining with amido black and Coomassie Blue | 44 |
| 3.2.9. | N-terminal and internal amino acid sequencing | 45 |
| 3.3. | Measurement of cytochrome c oxidase activity | 46 |
| 3.3.1. | Preparation of ferri- and ferro-cytochrome c solutions | 46 |
| 3.3.2. | Purification of asolectin and XAD-2 | 46 |
| 3.3.3. | Preparation of cytochrome c oxidase with regulatory properties | 47 |
| 3.3.4. | Reconstitution of cytochrome c oxidase for activity measurements | 47 |
| 3.3.5. | Phosphorylation and dephosphorylation of cytochrome c oxidase | 48 |
| 3.3.6. | Measurement of enzyme activity by the polarographical method | 48 |
| | Activity measurement of cytochrome c oxidase with regulatory properties | 49 |
| | Activity measurement of reconstituted cytochrome c oxidase | 49 |
| | Determination of respiratory control ratio (RCR) | 49 |
| 3.4. | Measurement of proton translocation | 50 |
| 3.4.1. | Reconstitution of cytochrome c oxidase for | |

| | |
|---|-----------|
| proton translocation measurements | 50 |
| 3.4.2. Proton translocation measurements using a pH electrode | 51 |
| 3.4.3. Proton translocation measurement using a stopped-flow spectrophotometer | 51 |
| 3.4.4. Proton conductivity measurement of proteoliposomes | 52 |
| 3.5. Immunochemical methods | 52 |
| Colorimetric detection of immunoreactivity in Western Blots using alkaline phosphatase | 52 |
| 4. Results | 54 |
| 4.1. Search for the cDNA of a new isoform of subunit IV (IV-2) | 54 |
| 4.1.1. Cloning and sequencing of subunit IV cDNA from rat | 56 |
| 4.1.2. Cloning and sequencing of subunit IV cDNA from turkey | 58 |
| 4.2. Search for cytochrome c oxidase subunit isoforms in bovine and turkey | 66 |
| 4.2.1. Comparison of bovine cytochrome c oxidase subunit patterns | 66 |
| 4.2.2. Searching for subunit IV-2 in bovine lung cytochrome c oxidase | 68 |
| 4.2.3. Comparison of subunit patterns of turkey cytochrome c oxidases | 71 |
| 4.2.4. Identification of subunit VIaL isoform in turkey liver, heart, and skeletal muscle | 73 |
| 4.3. Phosphorylation of cytochrome c oxidase subunits | 75 |
| 4.3.1. Identification of phosphorylated subunits of bovine heart cytochrome c oxidase with antibodies against phosphoserine and phosphothreonine | 75 |
| 4.3.2. Possible phosphorylation sites in the subunits of bovine heart cytochrome c oxidase | 77 |
| 4.4. Regulation of cytochrome c oxidase activity by cAMP-dependent phosphorylation | 81 |
| 4.4.1. The allosteric ATP-inhibition of cytochrome c oxidase is protected by KF and EGTA in the isolation media | 81 |
| 4.4.2. Effect of cAMP-dependent phosphorylation on the allosteric ATP-inhibition of cytochrome c oxidase isolated with KF and EGTA | 83 |
| 4.4.3. TMPD abolishes the allosteric ATP-inhibition of cytochrome c oxidase | 84 |
| 4.4.4. Incubation with protein phosphatase 1 releases the allosteric ATP-inhibition of cytochrome c oxidase | 85 |
| 4.4.5. Time dependent induction of allosteric ATP-inhibition of cytochrome c oxidase by cAMP-dependent phosphorylation | 86 |
| 4.4.6. Allosteric ATP-inhibition of cytochrome c oxidase is reversibly turned on by cAMP-dependent phosphorylation and turned off by dephosphorylation | 87 |
| 4.4.7. The allosteric ATP inhibition is obtained with cytochrome c oxidases from bovine heart, kidney, and testis | 88 |
| 4.4.8. Allosteric ATP-inhibition of cytochrome c oxidase is induced by phosphorylation from the cytosolic side | 90 |
| 4.4.9. Proposed phosphorylation site for allosteric ATP-inhibition | 91 |
| 4.5. Proton translocation in eukaryotic cytochrome c oxidase has a variable H ⁺ /e ⁻ stoichiometry | 94 |
| 4.5.1. Cardiolipin increases the H ⁺ /e ⁻ stoichiometry of reconstituted cytochrome c oxidase from bovine kidney but not from bovine heart | 94 |
| 4.5.2. Specific decrease of H ⁺ /e ⁻ stoichiometry by palmitate of cytochrome c oxidase from kidney but not from heart | 101 |
| 4.5.3. Palmitate increases the controlled respiration of kidney cytochrome c oxidase | 106 |

| | | |
|-----------|--|------------|
| 5. | Discussion | 110 |
| 5.1. | Search for cytochrome c oxidase subunit IV-2 isoform | 110 |
| 5.2. | Phosphorylation of Ser ⁴⁴¹ in subunit I of cytochrome c oxidase induces allosteric ATP-inhibition | 111 |
| 5.3. | Regulation of oxidative phosphorylation by allosteric ATP-inhibition of cytochrome c oxidase | 116 |
| 5.4. | Palmitate decreases the H ⁺ /e ⁻ stoichiometry of bovine kidney cytochrome c oxidase | 119 |
| 6. | Summary | 124 |
| 7. | Reference | 125 |
| 8. | Abbreviations | 139 |
| 9. | Acknowledgments | 141 |