



Rosemary Wanja Mutegi-Murori (Autor)

# Towards Identifying the Physiological and Molecular Basis of Drought Tolerance in Cassava (*Manihot esculenta* Crantz)

Department of Crop Sciences  
Georg-August University Göttingen



Rosemary Wanja Mutegi-Murori

---

Towards Identifying the Physiological and Molecular  
Basis of Drought Tolerance in Cassava  
(*Manihot esculenta* Crantz)

---



Cuvillier Verlag Göttingen  
Internationaler wissenschaftlicher Fachverlag

<https://cuvillier.de/de/shop/publications/851>

Copyright:

Cuvillier Verlag, Inhaberin Annette Jentzsch-Cuvillier, Nonnenstieg 8, 37075 Göttingen,  
Germany

Telefon: +49 (0)551 54724-0, E-Mail: [info@cuvillier.de](mailto:info@cuvillier.de), Website: <https://cuvillier.de>

## Table of Contents

<b>1. General Introduction .....</b>	<b>1</b>
1.1. Background .....	1
1.2. Importance of cassava.....	3
1.4. Biology of cassava .....	5
1.5. Cassava ecology and physiology .....	6
1.5.1. Sensitivity of cassava stomata .....	7
1.5.2. Leaf retention and changes in leaf expansion rates .....	7
1.5.3. Osmotic adjustment .....	8
1.5.4. Accumulation of specific low molecular weight proteins .....	8
1.5.5. Abscisic acid (ABA) accumulation .....	8
1.5.6. Accumulation and utilization of non-structural carbohydrate reserves	9
1.6. Mechanisms of drought tolerance in plants .....	9
1.6.1. Drought escape.....	10
1.6.2. Dehydration avoidance .....	10
1.6.3. Dehydration tolerance .....	10
1.7. Cassava breeding and biotechnology.....	11
1.8. Rationale of the study .....	12
1.9. Objectives of the study.....	13
1.10. Thesis outline .....	14
<b>2. Hardening of Cassava <i>In Vitro</i> Plantlets and Rapid Micro-propagation of Cassava Plants Through Nodal Cuttings.....</b>	<b>15</b>
2.1. Introduction.....	16
2.2. Materials and methods .....	19
2.2.1. Sub-culturing (in-direct micro-propagation).....	19
2.2.2. Transplanting and hardening (direct micro-propagation) .....	19
2.2.3. Rapid micro-propagation .....	20
2.3. Results and discussion .....	22
2.4. Conclusions.....	24
<b>3. Agronomic and Morphological Evaluation of Contrasting Cassava Germplasm Accessions under Drought Stress at Kiboko, Kenya .....</b>	<b>29</b>
3.1. Introduction.....	30
3.2. Materials and methods .....	33
3.2.1. Study site.....	33
3.2.2. Climate.....	33
3.2.3. Germplasm and field planting.....	34
3.2.4. Traits .....	36
3.2.5. Statistical analysis .....	37
3.3. Results.....	40
3.3.1. Plant height .....	40
3.3.2. Leaf retention .....	41
3.3.3. Number of branching levels.....	44
3.3.4. Leaf length and width .....	44
3.3.5. Harvest traits .....	46
3.3.6. Relationship between traits .....	46
3.3.7. Heritability .....	47

3.4. Discussion .....	51
3.5. Conclusion .....	55
<b>4. Metabolites Analysis in African Cassava Germplasm Accessions Evaluated at Kiboko Research Station, Makindu, Kenya.....</b>	<b>57</b>
4.1. Introduction.....	58
4.2. Materials and methods .....	61
4.2.1. Plant materials and sample preparation .....	61
4.2.2. Chromatography separation, ABA extraction and quantification.....	61
4.2.3. Determination of sugar content.....	63
4.2.4. Determination of protein content.....	64
4.2.5. Determination of starch content.....	64
4.2.6. Determination of amylose content in roots.....	64
4.2.7. Generation of derived variables .....	65
4.2.8. Statistical analysis .....	65
4.3. Results.....	67
4.3.1. Abscisic acid in cassava tissues .....	67
4.3.2. Assimilates content in cassava tissue.....	68
4.3.3. Relationship between traits .....	71
4.4. Discussion .....	72
<b>5. Genetic Mapping in Cassava (<i>Manihot esculenta</i> Crantz) using SSR's and EST-derived SSR's.....</b>	<b>79</b>
5.1. Introduction.....	80
5.2. Materials and methods .....	84
5.2.1. Mapping populations .....	84
5.2.1.1. Selection of contrasting parental accessions .....	84
5.2.1.2. Generation of crosses between contrasting parents .....	84
5.2.1.3. Segregating populations.....	86
5.2.2. Marker analysis.....	86
5.2.2.1. DNA isolation .....	86
5.2.2.2. Optimization of primers and labeling of ESSR's .....	87
5.2.2.3. Polymorphism screening and high throughput genotyping .....	88
5.2.2.4. Data scoring and linkage analysis.....	89
5.3. Results.....	92
5.3.1. Labeling of ESSR's, polymorphism screen and marker segregation ..	92
5.3.2. Map construction .....	93
5.3.3. Female and male map comparison.....	95
5.4. Discussion .....	97
5.4.1. Labeling of ESSR's, polymorphism screen and marker segregation ..	97
5.4.2. Map construction .....	100
5.4.3. Comparison with other maps .....	101
5.4.4. Female and male map comparison.....	102
<b>6. Summary.....</b>	<b>121</b>
<b>References.....</b>	<b>125</b>
<b>Acknowledgements .....</b>	<b>145</b>
<b>Curriculum Vitae .....</b>	<b>147</b>

## List of Tables

Table 1.1. World utilization patterns of cassava.	4
Table 2.1. Putative drought-tolerant and drought-susceptible African cassava germplasm accessions obtained from the IITA, Nigeria.	23
Table 2.2. Cost comparison of direct and in-direct methods of micro-propagating putative drought-tolerant and drought-susceptible cassava germplasm accessions.	25
Table 3.1. Physical soil characteristics of Kiboko Research Station, Makindu, Kenya.	33
Table 3.2. Putative drought-tolerant and drought-susceptible African cassava accessions utilized for field evaluation at Kiboko Research Station.	36
Table 3.3. Agronomic and morphological traits studied when evaluating putative drought-tolerant and drought-susceptible cassava accessions	39
Table 3.4. Responses to water stress of 31 cassava accessions at harvest	40
Table 3.5. Mean plant height of 31 cassava germplasm accessions evaluated at Kiboko Research Station, during different stress phases.	43
Table 3.6. Analysis of variance of 31 cassava accessions evaluated at Kiboko Research Station, Makindu, Kenya.	44
Table 3.7. Means of yield traits at harvest of 31 cassava germplasm accessions evaluated at Kiboko Research Station under well-watered and water-stressed treatments.	48
Table 3.8. Spearman's rank correlation coefficient between traits assessed for 31 cassava accessions evaluated at Kiboko Research Station, Makindu, Kenya.	49
Table 3.9. Spearman's rank correlation coefficient for individual treatments for 31 cassava accessions evaluated at Kiboko Research Station, Makindu, Kenya	50
Table 4.1. Analysis of variance of 31 cassava germplasm accessions evaluated at Kiboko Research Station.	68
Table 4.2. Spearman's rank correlation coefficient for the various metabolic traits evaluated for 31 cassava accessions at Kiboko Research Station.	71
Table 5.1. Population name, cross name and status of cassava germplasm accessions used in the generation of segregating populations.	85

Table 5.2. Polymerase Chain Reaction (PCR) optimization conditions developed by IITA-BecA, Nairobi.	88
Table 5.3. Properties of dyes used for fluorescent labeling of ESSR markers.	89
Table 5.4. Segregation types of markers mapped in the progeny of accessions.	93
Table 5.5. Details of the cassava female (COL 1734) and male (BRA 1149) genetic maps.	96
Table 5.6. Number of allelic bridges identified between the female and male genetic maps of cassava derived from a cross between a drought-tolerant and a drought-susceptible accession.	96