




Ingrid Maria Nijholt (Autor)

Molecular mechanisms of synaptic plasticity

Implications for immediate early gene expression and learning



<https://cuvillier.de/de/shop/publications/3571>

Copyright:

Cuvillier Verlag, Inhaberin Annette Jentsch-Cuvillier, Nonnenstieg 8, 37075 Göttingen, Germany

Telefon: +49 (0)551 54724-0, E-Mail: info@cuvillier.de, Website: <https://cuvillier.de>

CONTENTS

| | page |
|---|---------|
| <u>Chapter 1:</u> General introduction | 1 |
| 1. Synaptic plasticity | 1 |
| 1.1 Basic principles of synaptic transmission | 2 |
| 1.2 Second messenger pathways | 3 |
| 1.2.1 Protein kinases | 4 |
| 1.2.2 Protein phosphatases | 5 |
| 1.2.3 Phosphoproteins | 6 |
| 1.3 Glutamate receptors | 7 |
| 1.3.1 NMDA receptor | 7 |
| 2. Synaptic plasticity and gene expression | 9 |
| 2.1 CREB | 10 |
| 2.2 FOS | 11 |
| 3. Synaptic plasticity and learning and memory | 12 |
| 3.1 Hippocampus | 12 |
| 3.2 Long-term potentiation and depression as a model for learning and memory | 12 |
| 3.3 Influence of stress on synaptic plasticity and learning and memory | 14 |
| 3.3.1 Corticotropin-releasing factor as mediator of the stress response | 15 |
| <u>Chapter 2:</u> Modulation of hypothalamic NMDA receptor function by cyclic AMP-dependent protein kinase and phosphatases | 33 |
| <u>Chapter 3:</u> The phosphoprotein DARPP-32 mediates cAMP-dependent potentiation of striatal N-methyl-D-aspartate responses | 45 |
| <u>Chapter 4:</u> <i>In vivo</i> NMDA/dopamine interaction resulting in FOS production in the limbic system and basal ganglia of the mouse brain. | 62 |
| <u>Chapter 5:</u> <i>In vivo</i> CREB phosphorylation mediated by dopamine and NMDA receptor activation in mouse hippocampus and caudate nucleus. | 79 |
| <u>Chapter 6:</u> The corticotropin-releasing factor receptor 1 antagonist CP-154,526 markedly enhances associative learning and paired-pulse facilitation immediately after a stressful experience. | 87 |
| <u>Chapter 7:</u> Priming of long-term potentiation in mouse hippocampus by corticotropin-releasing factor and acute stress: implications for hippocampus-dependent learning. | 101 |

| | |
|--|-----|
| <u>Chapter 8: Modulation of neuronal excitability and associative learning by different corticotropin-releasing factor-signaling cascades in mouse hippocampus</u> | 117 |
| <u>Chapter 9: Summary and general discussion</u> | 135 |
| 1. Brain regional differences in the modulation of the NMDA receptors by phosphorylation events | 135 |
| 1.1 DARPP-32 | 136 |
| 1.2 Possible physiological relevance of PKA-mediated NMDA receptor modulation in the hypothalamus | 138 |
| 1.3 Direct versus indirect phosphorylation of the NMDA receptor | 138 |
| 2. Brain regional differences in gene expression | 139 |
| 2.1 Specificity in signaling pathways | 140 |
| 3. Synaptic plasticity and learning and memory | 143 |
| 3.1 Metaplasticity | 143 |
| 3.2 Correlation between synaptic plasticity and learning | 146 |
| 3.2.1 Possible role of calcium/calmodulin-dependent protein kinase II | 147 |
| 3.2.2 Comparison between two mouse inbred strains Balb/c and C57BL/6N | 148 |
| 4. Concluding remarks | 149 |
| <u>Nederlandse samenvatting</u> | 159 |
| <u>Supplement: NS-257, a novel competitive AMPA receptor antagonist, interacts with kainate and NMDA receptors.</u> | 165 |
| <u>List of publications</u> | 179 |
| <u>Dankwoord</u> | 183 |
| <u>Curriculum vitae</u> | 185 |