



# Table of contents

<b>Chapter 1 Introduction .....</b>	<b>1</b>
1.1 Molecular magnets .....	1
1.2 Definition of molecular magnetism .....	3
1.2.1 Magnetic materials .....	3
1.2.2 The description of the magnetic behavior .....	4
1.3 Single molecule magnets .....	8
1.4 Fe <sup>III</sup> -Ln <sup>III</sup> SMMs .....	12
1.5 <sup>57</sup> Fe Mössbauer spectroscopy .....	14
1.5.1 Fundamental of Mössbauer spectroscopy .....	14
1.5.2 Mössbauer parameters .....	15
1.5.3 Mössbauer spectroscopy for Fe <sup>III</sup> -Ln <sup>III</sup> coordination clusters .....	17
1.6 Synthetic Strategies to Fe <sup>III/II</sup> -Ln <sup>III</sup> coordination clusters .....	18
1.7 Ligand Selection .....	20
1.8 Research objectives .....	22
1.9 Thesis overview .....	24
<b>Chapter 2 Fe<sup>III</sup>-Ln<sup>III</sup> coordination clusters with diethylene glycol (H<sub>2</sub>deg) as ligand .....</b>	<b>25</b>
2.1 Introduction .....	25
2.2 Synthetic approach .....	26
2.3.1 Structures of [HN(C <sub>2</sub> H <sub>5</sub> ) <sub>3</sub> ][Fe <sup>III</sup> <sub>4</sub> Ln <sup>III</sup> <sub>3</sub> (μ <sub>3</sub> -OH) <sub>2</sub> (deg) <sub>5</sub> (N <sub>3</sub> ) <sub>6</sub> (ben) <sub>4</sub> ] ·xCH <sub>3</sub> CN·yH <sub>2</sub> deg·zH <sub>2</sub> O [Ln = Y( <b>1</b> ), Tb( <b>2</b> ), Dy( <b>3</b> ), Ho( <b>4</b> ); x = 0.7, y = 1, z = 0 for ( <b>1</b> ); x = 1.3, y = 0.4, z = 3.7 for ( <b>2</b> ); x = 0.1, y = 0.8, z = 0 for ( <b>3</b> ); x = 0.3, y = 1, z = 0 for ( <b>4</b> )] .....	27
2.3.2 Magnetic properties of compounds <b>1-4</b> .....	29
2.3.3 Mössbauer spectra for compounds <b>1-3</b> .....	34
2.4.1 Structures of [Fe <sup>III</sup> <sub>2</sub> Ln <sup>III</sup> <sub>3</sub> (μ <sub>3</sub> -OH) <sub>6</sub> (H <sub>2</sub> deg) <sub>3</sub> (ben) <sub>6</sub> ]Cl <sub>3</sub> ·xCH <sub>3</sub> CN·yH <sub>2</sub> O [Ln = Y( <b>5</b> ), Sm( <b>6</b> ), Eu( <b>7</b> ), Gd( <b>8</b> ), Tb( <b>9</b> ), Dy( <b>10</b> ), Ho( <b>11</b> ); x = 0.6, y = 3.5 for ( <b>5</b> ), ( <b>11</b> ); x = 1.8, y = 2 for ( <b>6</b> ); x = 0.5, y = 2 for ( <b>7</b> ); x = 1, y = 2 for ( <b>8</b> ); x = 1.5, y = 2 for ( <b>9</b> ); x = 1.7, y = 2 for ( <b>10</b> )] .....	39
2.4.2 Magnetic properties of compounds <b>5-11</b> .....	41
2.4.3 Structures of [Fe <sup>III</sup> <sub>2</sub> Ln <sup>III</sup> <sub>3</sub> (μ <sub>3</sub> -OH) <sub>6</sub> (H <sub>2</sub> deg) <sub>3</sub> (naph) <sub>6</sub> ]Cl <sub>3</sub> ·xCH <sub>3</sub> CN·yH <sub>2</sub> O [Ln = Y( <b>12</b> ), Tb( <b>13</b> ), Dy( <b>14</b> ), Ho( <b>15</b> ); x = 1.8, y = 4.5 for ( <b>12</b> ); x = 0.5, y = 3 for ( <b>13</b> ); x = 1.7, y = 3 for ( <b>14</b> ); x = 1.8, y = 2.8 for ( <b>15</b> )] .....	47

2.4.4 Magnetic properties of compounds <b>12-15</b> .....	48
2.5.1 Structures of $[Fe^{III}{}_4Ln^{III}{}_4(\mu_4\text{-O})_2(\mu_3\text{-OH})_4(\mu_2\text{-OH})(Hdeg)_2(\text{ben})_{12}]Cl \cdot xCH_3CN \cdot yH_2O$ .....	53
$[Ln = Y(\mathbf{16}), Tb(\mathbf{17}), Dy(\mathbf{18}), Ho(\mathbf{19}), Er(\mathbf{20}), Tm(\mathbf{21}), Yb(\mathbf{22}); x = 1.3, y = 7 \text{ for } (\mathbf{16}), (\mathbf{19}), (\mathbf{22}); x = 1, y = 4.5 \text{ for } (\mathbf{17}); x = 1.1, y = 7 \text{ for } (\mathbf{18}); x = 1.4, y = 7 \text{ for } (\mathbf{20}); x = 1.5, y = 7 \text{ for } (\mathbf{21})]$ .....	53
2.5.2 Magnetic properties of compounds <b>16-22</b> .....	55
2.5.3 Mössbauer spectra of compounds <b>16</b> and <b>18</b> .....	59
<b>Chapter 3 Fe<sup>III</sup>-Ln<sup>III</sup> coordination clusters with triethylene glycol (H<sub>2</sub>teg) as ligand</b> .....	<b>62</b>
3.1 Introduction.....	62
3.2 Synthetic approach.....	62
3.3.1 Structures of $[Fe^{III}{}_2Ln^{III}{}_2(\mu_3\text{-OH})_2(teg)_2(N_3)_{2-x}Cl_x(piv)_4]$ [ $Ln = Y(\mathbf{23}), Dy(\mathbf{24})$ and $Ho(\mathbf{25})$ ; $x = 0.2$ for $(\mathbf{23})$ ; $x = 0$ for $(\mathbf{24}), (\mathbf{25})$ ].....	64
3.3.2 Structures of $[Fe^{III}{}_2Ln^{III}{}_2(\mu_3\text{-MeO})_2(teg)_2(NO_3)_2(piv)_4] \cdot xH_2O$ [ $Ln = Y (\mathbf{26}), Dy (\mathbf{27})$ and $Ho (\mathbf{28})$ ; $x=0$ for $(\mathbf{26})$ ; $x= 1$ for $(\mathbf{27}), (\mathbf{28})$ ] .....	66
3.3.3 Magnetic properties of <b>23-28</b> .....	67
3.3.4 The structure of $[Fe^{III}{}_2Ln^{III}{}_2(\mu_3\text{-OH})_2(teg)_2(N_3)_{2-x}Cl_x(\text{ben})_4]$ [ $Ln = Y (\mathbf{29a}), Dy (\mathbf{29b})$ ; $x = 0.2$ for $(\mathbf{29a})$ ; $x = 0$ for $(\mathbf{29b})$ ].....	71
3.3.5 Magnetic properties of <b>29a</b> and <b>29b</b> .....	72
3.3.6 Mössbauer spectra of compounds <b>29a</b> and <b>29b</b> .....	75
3.4.1 Structures of $[Fe^{III}{}_2Ln^{III}{}_3(\mu_4\text{-O})(\mu_3\text{-OH})(teg)_2(N_3)(NO_3)(piv)_6]$ [ $Ln = Tb(\mathbf{30}), Dy(\mathbf{31}), Ho(\mathbf{32})$ ].....	79
3.4.2 Structures of $[Fe^{III}{}_2Ln^{III}{}_3(\mu_4\text{-O})(\mu_3\text{-OH})(teg)_2(piv)_8] \cdot xCH_3CN$ [ $Ln = Y(\mathbf{33}), Sm(\mathbf{34}), Eu(\mathbf{35}), Gd(\mathbf{36}), Tb(\mathbf{37}), Dy(\mathbf{38}), Ho(\mathbf{39}), Er(\mathbf{40}), Tm(\mathbf{41})$ ; $x = 0$ for $(\mathbf{33}), (\mathbf{37})-(\mathbf{41})$ ; $x = 2$ for $(\mathbf{34})$ ; $x = 1.5$ for $(\mathbf{35})$ ; $x = 1$ for $(\mathbf{36})$ ] .....	81
3.4.3 Magnetic properties of compounds <b>30-32</b> .....	83
3.4.4 Magnetic properties of compounds <b>33-41</b> .....	87
3.4.5 Mössbauer spectra of compounds <b>31, 33</b> and <b>38</b> .....	94
3.5.1 Structures of $[Fe^{III}{}_5Ln^{III}{}_2(\mu_4\text{-O})_2(\mu_3\text{-O})(\mu_3\text{-OH})(teg)(piv)_{12}] \cdot xH_2O$ [ $Ln = Y(\mathbf{42}), Tb(\mathbf{43}), Dy(\mathbf{44}), Ho(\mathbf{45}), Er(\mathbf{46}), Tm(\mathbf{47}), Yb(\mathbf{48})$ ; $x = 0$ for $(\mathbf{42})-(\mathbf{46})$ ; $x = 1$ for $(\mathbf{47})$ ; $x = 0.5$ for $(\mathbf{48})$ ].....	97
3.5.2 Magnetic properties of compounds <b>42-48</b> .....	99
3.5.3 Mössbauer spectra of compounds <b>42</b> and <b>44</b> .....	106
<b>Chapter 4 Fe<sup>III</sup>-Ln<sup>III</sup> coordination clusters with tetraethylene glycol (H<sub>2</sub>taeg) as ligand</b> ... <b>111</b>	
4.1 Introduction.....	111
4.2 Synthesis .....	112
4.3.1 Structure of $[Fe^{III}{}_3Ln^{III}{}_4(\mu_3\text{-OH})_6(taeg)_3(piv)_6Cl]Cl \cdot 0.5[Ln(NO_3)_5] \cdot xCH_3CN$ [ $Ln = Gd (\mathbf{49}), Tb(\mathbf{50}), Dy(\mathbf{51}), Ho(\mathbf{52})$ ; $x= 2$ for $(\mathbf{49})$ ; $x= 1.4$ for $(\mathbf{50}), (\mathbf{51})$ ; $x= 1.1$ for $(\mathbf{52})$ ].....	112
4.3.2 Magnetic properties of <b>49-52</b> .....	115



<b>Chapter 5 Conclusions .....</b>	<b>118</b>
<b>Chapter 6 Experimental section .....</b>	<b>126</b>
<b>Chapter 7 Crystallographic data.....</b>	<b>146</b>
<b>Chapter 8 Bibliography .....</b>	<b>153</b>
<b>Appendix .....</b>	<b>162</b>
<b>Acknowledgements .....</b>	<b>166</b>
<b>Curriculum Vitae .....</b>	<b>168</b>